

# Breast Cancer Screening (BCS)

Effectiveness of Care HEDIS® Measure

## Measure definition

Female patients ages 52–74 who had a mammogram to screen for breast cancer any time on or between October 1 two years prior to the measurement year and December 31 of the measurement year

## Exclusions

Patients are excluded if they:

- Have a history of mastectomy on both the left and right side on the same or different dates of service.
- Received hospice care during the measurement year.
- Are age 66 and older with advanced illness and frailty (for additional definition information, see the *Advanced Illness and Frailty Guide*).
- Are deceased during the measurement year.
- Received palliative care during the measurement year

## Information that patient medical records should include

- Date the mammogram was performed.
- Documentation of mastectomy and date performed (if exact date is unknown, the year is acceptable).

## Information that patient claims should include

If the patient met exclusion criteria, include the following ICD-10-CM<sup>1</sup> diagnosis codes on the claim, as appropriate:

ICD-10-CM code	Description
Z90.11	Acquired absence of right breast and nipple
Z90.12	Acquired absence of left breast and nipple
Z90.13	Acquired absence of bilateral breasts and nipples

## Tips for success

- Create a standing order to mail to patient for mammography.
- Provide a list of locations where mammogram screenings can be performed.
- If telehealth, telephone or e-visits are used instead of face-to-face visits, discuss the need for breast cancer screening and mail a mammogram order with location of testing facility and phone number.

## Tips for talking with patients

Educate patients about the importance of routine screening:

- Many women with breast cancer do not have symptoms, which is why regular breast cancer screenings are so important.
- Mammograms are an effective method for detecting breast cancer in early stages, when it is most treatable.<sup>2</sup>
- The recommended frequency of routine mammograms is at least once every 24 months for all women ages 50–74. Depending on risk factors, mammograms may be done more frequently.
- The accuracy of mammography improves as women age and has an overall detection rate of about 87 percent.<sup>3</sup>

<sup>1</sup>ICD-10-CM created by the National Center for Health Statistics, under authorization by the World Health Organization. WHO-copyright holder.

<sup>2</sup>American Cancer Society. <https://www.cancer.org/cancer/breast-cancer/screening-tests-and-early-detection/american-cancer-society-recommendations-for-the-early-detection-of-breast-cancer.html>

<sup>3</sup>Komen. <https://www.komen.org/breast-cancer/screening/mammography/accuracy/>