

Medical Policy



Title: Continuous Passive Motion in the Home Setting

Professional / Institutional
Original Effective Date: June 10, 1986 / March 8, 2010
Latest Review Date: April 23, 2024
Current Effective Date: June 7, 2021

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: • With total knee arthroplasty	Interventions of interest are: • Continuous passive motion in the home setting	Comparators of interest are: • Physical therapy alone • Standard of care, if unable to tolerate physical therapy	Relevant outcomes include: • Symptoms • Functional outcomes
Individuals: • With intra-articular cartilage repair of the knee	Interventions of interest are: • Continuous passive motion in the home setting	Comparators of interest are: • Standard of care	Relevant outcomes include: • Symptoms • Functional outcomes
Individuals: • With musculoskeletal conditions other than total knee arthroplasty	Interventions of interest are:	Comparators of interest are: • Standard of care	Relevant outcomes include: • Symptoms

Populations	Interventions	Comparators	Outcomes
or knee cartilage repair requiring physical therapy	<ul style="list-style-type: none"> • Continuous passive motion in the home setting 		<ul style="list-style-type: none"> • Functional outcomes
Individuals: <ul style="list-style-type: none"> • Who have had a stroke requiring physical therapy 	Interventions of interest are: <ul style="list-style-type: none"> • Continuous passive motion in the home setting 	Comparators of interest are: <ul style="list-style-type: none"> • Standard of care 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Functional outcomes

DESCRIPTION

Continuous passive motion devices are used to keep a joint in motion without patient assistance. Continuous passive motion is being evaluated for treatment and postsurgical rehabilitation of the upper- and lower-limb joints and for a variety of musculoskeletal conditions.

OBJECTIVE

The objective of this evidence review is to determine whether continuous passive motion improves the net health outcome in individuals undergoing postsurgical rehabilitation in the home setting

BACKGROUND

Physical therapy of joints following surgery focuses both on passive motion to restore mobility and on active exercises to restore strength. While passive motion can be administered by a therapist, continuous passive motion devices have also been used. Continuous passive motion is thought to improve recovery by stimulating the healing of articular tissues and the circulation of synovial fluid; reducing local edema; and preventing adhesions, joint stiffness or contractures, or cartilage degeneration.¹ Continuous passive motion has been investigated primarily in the knee, particularly after total knee arthroplasty or ligamentous or cartilage repair. Acceptance of its use in the knee joint has created interest in continuous passive motion use for other weight-bearing joints (i.e., hip, ankle, metatarsals) as well as non-weight-bearing joints (i.e., shoulder, elbow, metacarpals, interphalangeal joints). Use of continuous passive motion in stroke and burn patients is also being explored.

The device used for the knee moves the joint (e.g., flexion and extension) without patient assistance, continuously for extended periods of time (i.e., up to 24 hours/day).¹ An electrical power unit is used to set the variable range of motion and speed. The initial settings for range of motion are based on a patient’s level of comfort and other factors assessed intraoperatively. The range of motion is increased by 3 to 5 degrees per day, as tolerated. The speed and range of motion can be varied, depending on joint stability. The use of the device may be initiated in the immediate postoperative period and then continued at home for a variable period of time.

Over time, hospital lengths of stay have progressively shortened and, in some cases, surgical repair is done as an outpatient or with a length of stay of 1 to 2 days.² As a result, there has been a considerable shift in the rehabilitation regimen, moving range of motion from an intensive in-hospital program to a less intensive outpatient program. Some providers may want patients to

continue continuous passive motion in the home setting as a means of duplicating services offered with a longer (7-day) hospital stay.

The focus of the current review is to examine the literature on the use of continuous passive motion in the home setting as it is currently being prescribed postoperatively. Relevant comparisons are treatment outcomes of continuous passive motion when used alone or with physical therapy, compared with physical therapy alone.

REGULATORY STATUS

Continuous passive motion devices are considered class 1 devices by the U.S. Food and Drug Administration and are exempt from 510(k) requirements. This classification does not require submission of clinical data on efficacy but only notification of the Food and Drug Administration prior to marketing. Food and Drug Administration product code: BXB.

POLICY

- A. Use of continuous passive motion (CPM) in the home setting may be considered **medically necessary** as an adjunct to physical therapy in the following situations:
1. Under conditions of low postoperative mobility or inability to comply with rehabilitation exercises following a total knee arthroplasty (TKA) or TKA revision. This may include individuals with complex regional pain syndrome (reflex sympathetic dystrophy); extensive arthrofibrosis or tendon fibrosis; or physical, mental, or behavioral inability to participate in active physical therapy.
 2. During the non-weight-bearing rehabilitation period following intra-articular cartilage repair procedures of the knee (e.g., microfracture, osteochondral grafting, autologous chondrocyte implantation, treatment of osteochondritis dissecans, repair of tibial plateau fractures).
- B. Use of CPM in the home setting for all other conditions is considered **not medically necessary**.

POLICY GUIDELINES

- A. Following total knee arthroplasty, continuous passive motion in the home setting will be allowable for up to 17 days after surgery while individuals are immobile or unable to bear weight.
- B. Following articular cartilage repair procedures of the knee, continuous passive motion in the home setting will be allowable for up to 6 weeks during non-weight-bearing rehabilitation.

Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through January 17, 2024.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, two domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

TOTAL KNEE ARTHROPLASTY

Clinical Context and Therapy Purpose

The purpose of continuous passive motion in the home setting in individuals with total knee arthroplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review:

Populations

The relevant population of interest is individuals with total knee arthroplasty.

Interventions

The therapy being considered is continuous passive motion.

Comparators

The following therapies are currently being used for total knee arthroplasty: physical therapy (PT) alone or standard of care, if unable to tolerate PT.

Outcomes

The general outcomes of interest are symptoms and functional outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;

- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

EARLY POSTOPERATIVE IN-HOSPITAL SETTING

REVIEW OF EVIDENCE

Systematic Reviews

This evidence review was informed by a TEC Assessment (1997) that concluded continuous passive motion met the TEC criteria as an adjunct to PT in patients undergoing total knee arthroplasty.³ Early studies of continuous passive motion machines focused on their use in the hospital setting, in which the impact on length of stay was frequently considered a key clinical outcome, and so the TEC Assessment did not specifically examine the point of service or the length of time continuous passive motion devices were used. A critical study identified in the TEC Assessment was an RCT by McInnes et al (1992) that compared use of continuous passive motion initiated in the immediate postoperative period and continued through the 7-day hospital stay with standard rehabilitation alone.⁴ At 6 weeks postoperatively, the most salient difference between groups was an increased incidence of arthrofibrosis requiring manipulation in the non-continuous passive motion group.

Efficacy in the early postoperative period has been cited as a reason to support the continued use of these devices in the non-acute care hospital or home setting following early discharge. Continuous passive motion after total knee arthroplasty was the subject of a 2003 Cochrane review.⁵ Reviewers reported that continuous passive motion combined with PT significantly increased active knee flexion and decreased length of stay. However, the analysis suggested the benefits of continuous passive motion in a hospital setting may be small and only short-term.⁶ This Cochrane review was updated in 2010 and again in 2014.^{7,8} The updated review included 24 RCTs with 1445 participants and examined short-term (<6 weeks), medium-term (6 weeks to 6 months), and long-term (>6 months) effects of continuous passive motion. Most selected studies examined short-term effects. Continuous passive motion was applied for 1.5 to 24 hours a day, over 1 to 17 days. A summary of findings is provided in Table 1.

Table 1. 2014 Cochrane Review Findings on Continuous Passive Motion

Findings	QOE
continuous passive motion increases passive and active knee flexion range of motion (mean difference, 2), but the effects were too small to be clinically relevant	Moderate
continuous passive motion does not have clinically important short-term effects on pain (-0.4 points on a 10-point scale)	Low
continuous passive motion does not have clinically important medium-term effects on function or quality of life	Moderate

Findings	QOE
continuous passive motion may reduce the need for manipulation under anesthesia (25 fewer manipulations per 1000; RR , 0.3)	Very low
continuous passive motion reduced the risk of adverse events (13 fewer adverse events per 1000 ; RR , 0.9)	Low

Adapted from Harvey et al (2014).⁷ QOE: quality of evidence; RR: relative risk.

Another 2014 Cochrane systematic review, which included 11 RCTs, found no evidence that continuous passive motion reduced venous thromboembolism after total knee arthroplasty.⁹

Randomized Controlled Trials

Yashar et al (1997) randomized 178 patients undergoing total knee arthroplasty to continuous passive motion immediately postsurgery or to continuous passive motion 1 day postsurgery.¹⁰ A small but statistically significant improvement in flexion was found at the time of discharge among those started on immediate continuous passive motion, but this difference did not persist at 4 weeks. MacDonald et al (2000) reported on a randomized trial comparing immediate postoperative continuous passive motion with no continuous passive motion for 120 patients after total knee arthroplasty.¹¹ Patients received a maximum of 24 hours with continuous passive motion. There were no differences between treatment groups in range of motion, length of stay, or analgesic requirements. In a trial reported by Pope et al (1997), 53 patients were randomized to 1 of 2 schedules of continuous passive motion (both for 48 hours) or to no continuous passive motion.¹² The use of continuous passive motion was not associated with improved long-term function or range of motion. Kumar et al (1996) randomized 73 patients who had undergone total knee arthroplasty to continuous passive motion immediately postsurgery or to a protocol of early passive flexion, referred to as the "drop and dangle" technique.¹³ Patients assigned to passive flexion were discharged from the hospital one day earlier and also had a statistically better extension range of 2.8 at 6 months than the continuous passive motion group.

Other RCTs have found that 2 to 4 hours of daily continuous passive motion in the hospital after total knee arthroplasty did not improve postoperative outcomes at discharge or follow-up.^{14,15,16,17} In one trial, Bruun-Olsen et al (2009) randomized 63 patients undergoing total knee arthroplasty to active PT exercises with or without continuous passive motion to assess any short-term benefit on pain or function.¹⁴ In both groups, exercises were performed daily for 30 minutes, starting 1 day after surgery and continuing until discharge at 1 week. For the experimental group, continuous passive motion was administered for 4 hours on the day of surgery, followed by 6 hours daily in addition to therapist-guided exercises. Blinded assessments at 1 week and 3 months after surgery showed similar results for pain and function in the 2 groups. At 1 week, both groups had visual analog scale pain ratings of 40 and flexion scores within 2 of each other. Functional testing at 3 months showed no benefit of adjunctive continuous passive motion. The lack of improvement with continuous passive motion in these studies might have been attributable to patients mobilizing or commencing flexion immediately following surgery.¹⁶ A 2014 study of 150 patients undergoing total knee arthroplasty found no benefit of continuous passive motion when used over a 2-day postoperative hospital stay.¹⁷

Non-Acute Care Hospital Setting

In a RCT, Herbold et al (2014) assessed 141 total knee arthroplasty patients assigned to daily conventional therapy lasting 3 hours or daily continuous passive motion for 2 hours throughout

their inpatient rehabilitation stay.¹⁸ After an average length of stay of 8 days, there were no significant differences between the continuous passive motion and no continuous passive motion groups for active range of motion, Timed Up and Go test, knee girth, Functional Independence Measure scores, ambulation device at discharge, or on the self-reported Western Ontario and McMaster Universities Osteoarthritis Index scores.

In 2000, Chen et al (2000) randomized 51 patients in an inpatient rehabilitation service who had undergone total knee arthroplasty to conventional active PT or to PT plus continuous passive motion.¹⁹ Referral to the rehabilitation center was made 5 to 6 days after surgery, and most had received continuous passive motion as part of the initial hospitalization. Knee flexion was the principal outcome. No significant differences were noted in passive range of motion between the 2 groups, as measured on admission, on the third and seventh days, and at the time of discharge (8 days postadmission). Thus, the use of continuous passive motion in the rehabilitation hospital offered no added benefit.

In a 2012 retrospective comparative study, the same group as the Herbold et al RCT evaluated the use of continuous passive motion in 61 matched pairs of patients admitted to a rehabilitation hospital.²⁰ Outcomes following use of continuous passive motion were compared with those from a cohort of 61 inpatients who also had poor initial range of motion, defined as less than 75° of active knee flexion at the time of admission, and matched for postoperative day at admission, age, length of stay, and Health Insurance Prospective Payment System code. Use of continuous passive motion (2 hours/day) was determined primarily by the referring physician and used in 29% of the pool of 633 patients who had poor initial range of motion. Average length of stay was 7.85 days. There were no significant differences in outcomes at discharge, including knee flexion or extension, discharge to the community, need for home care services, need for an assistive device, or functional scores on the Health Insurance Prospective Payment System.

Home Setting

A study by Worland et al (1998) compared the use of continuous passive motion with active PT in the home setting. At discharge, they randomized 80 patients undergoing total knee arthroplasty to home continuous passive motion (3 hours/day for 10 days) or to active PT.²¹ Most studies have examined continuous passive motion as an adjunct to active PT, while this study proposed continuous passive motion as an alternative to PT. At 2 weeks, knee flexion was similar in both groups but a flexion contracture was noted in 1 patient in the continuous passive motion group. At 6 months, no differences were found in knee scores or knee flexion.

In another RCT, Lenssen et al (2008) evaluated 60 patients with limited flexion range of motion (<80°) at the time of hospital discharge who were assigned to standard PT alone or PT plus continuous passive motion in the home (4 hours/day) until assessment on postoperative day 17.²² Blinded assessment showed a trend for increased range of motion for the continuous passive motion group (e.g., 89 vs. 84, respectively ; $p=.07$), with no differences in function between groups, as measured by the Knee Society Score (function subscore 43 vs. 40, respectively) and the Western Ontario and McMaster Universities Osteoarthritis Index difficulty score (49 vs. 45, respectively). No differences were observed between groups in range of motion or function at the 6-week or 3-month assessment. In addition, no differences were observed for the secondary outcome measures (perceived effect, medication use, satisfaction with treatment, adherence) at either of the assessment times.

Section Summary: Total Knee Arthroplasty

Numerous RCTs have compared continuous passive motion as adjunctive therapy with PT for patients undergoing total knee arthroplasty. Most trials used continuous passive motion in the inpatient setting and are less relevant to today's practice patterns of shorter hospital stays followed by outpatient rehabilitation. Some of these trials reported improvements in range of motion for patients receiving continuous passive motion but these improvements were short-term, of small magnitude, and of uncertain clinical significance. The RCTs that specifically evaluated continuous passive motion in the non-acute care hospital setting or home setting did not show improved outcomes with continuous passive motion.

ARTICULAR CARTILAGE REPAIR OF THE KNEE**Clinical Context and Therapy Purpose**

The purpose of continuous passive motion in the home setting in individuals with articular cartilage repair of the knee is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review:

Populations

The relevant population of interest is individuals with articular cartilage repair of the knee.

Interventions

The therapy being considered is continuous passive motion.

Comparators

The following therapies are currently being used for articular cartilage repair of the knee: standard of care.

Outcomes

The general outcomes of interest are symptoms and functional outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Although no RCTs were identified comparing health outcomes with or without the use of continuous passive motion, continuous passive motion is routinely used as part of the rehabilitation protocol for as long as 6 weeks when weight-bearing is restricted following autologous chondrocyte implantation.^{23,24,25} Basic research supports the use of continuous passive motion to facilitate greater healing of articular cartilage of full-thickness defects that penetrate the subchondral bone compared with either immobilization or intermittent mobilization.^{26,27}

Fazalare et al (2010) published a systematic review of continuous passive motion after knee cartilage defect surgery.²⁸ Reviewers found that continuous passive motion had been used following autologous chondrocyte implantation, microfracture, and osteochondral autografts in numerous studies in the previous 5 years. Four level III (cohort) studies with 262 patients were identified that compared continuous passive motion with no continuous passive motion; no RCTs were identified. Procedures in these 4 studies included microfracture, periosteal transplant of the patella, and high tibial osteotomy with diagnostic arthroscopy or abrasion arthroplasty. Continuous passive motion regimens ranged from 6 days to 8 weeks. Heterogeneity in the studies and outdated surgical techniques limited conclusions drawn from these trials. Clinical outcomes did not permit a definitive conclusion of efficacy of continuous passive motion. However, reviewers cited several studies in which other outcomes (e.g., histologic outcomes on follow-up biopsies) did favor continuous passive motion.

Another systematic review by Howard et al (2010) evaluated continuous passive motion and other postoperative practices after knee cartilage repair.²⁹ Reviewers cited several basic science studies using animal models that appear to support continuous passive motion. They identified 2 clinical studies, both of which were retrospective nonrandomized comparative studies. In 1 study (N =43), there were no differences between groups in clinical or functional outcomes at an average follow-up of 4.2 years. In the other study (N =77), patients in the continuous passive motion group (n=46) had greater improvement in grading of the cartilage lesion compared with patients who did not have access to continuous passive motion (n=31).

Section Summary: Articular Cartilage Repair of the Knee

Current evidence on use of continuous passive motion to facilitate knee rehabilitation after articular cartilage repair includes systematic reviews. These reviews reported methodologic issues with available cohort studies and a paucity of studies assessing clinical application of continuous passive motion to knee rehabilitation.

OTHER MUSCULOSKELETAL CONDITIONS REQUIRING PHYSICAL THERAPY

Clinical Context and Therapy Purpose

The purpose of continuous passive motion in the home setting in individuals with musculoskeletal conditions other than total knee arthroplasty or knee cartilage repair requiring PT is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review:

Populations

The relevant population of interest is individuals with musculoskeletal conditions other than total knee arthroplasty or knee cartilage repair requiring PT.

Interventions

The therapy being considered is continuous passive motion.

Comparators

The following therapies are currently being used for musculoskeletal conditions other than total knee arthroplasty or knee cartilage repair requiring PT: standard of care.

Outcomes

The general outcomes of interest are symptoms and functional outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE**Articular Knee Fractures**

Hill et al (2014) randomized 40 patients with articular fractures of either the proximal part of the tibia or the distal end of the femur to standardized PT with or without continuous passive motion for 48 hours postoperatively.³⁰ At the 48-hour assessment, the continuous passive motion group had significantly greater knee flexion (43 difference ; $p < .005$). However, 6 of 20 patients were unable to tolerate continuous passive motion and there was no benefit to adding 48 hours of continuous passive motion when assessed at any of the follow-up visits (2, 6, 12, and 24 weeks).

Anterior Cruciate Ligament Repair

This literature review did not identify any RCTs of continuous passive motion in the home setting after repair of the anterior cruciate ligament. However, the studies of continuous passive motion after anterior cruciate ligament repair in the immediate postoperative period may be relevant to the non-acute care hospital or home setting for patients discharged following a shorter hospital stay. The TEC Assessment (1997) concluded that continuous passive motion as an adjunct to conventional PT in the immediate postoperative period after anterior cruciate ligament repair offered no demonstrable advantage over conventional PT alone.³ In a systematic review of anterior cruciate ligament reconstruction rehabilitation, Wright et al (2008) discussed 6 RCTs on continuous passive motion published before 1996; no RCTs published after the 1997 TEC Assessment were identified.³¹ Reviewers found no substantial advantage for continuous passive

motion use and concluded that continuous passive motion for anterior cruciate ligament rehabilitation could not be justified. Wright et al (2008) also noted that most current anterior cruciate ligament rehabilitation protocols initiate early motion within the first postoperative week.

A 2022 review was conducted to synthesize evidence from systematic reviews for rehabilitation interventions following anterior cruciate ligament injury.³² This review identified 1 systematic review that included evidence for continuous passive motion by Gatewood et al (2017).³³ The authors identified 2 RCTs of continuous passive motion in the immediate postoperative setting, 1 of which was not included in the review by Wright et al (2008). In this study, 60 patients (95% of whom were men) were randomized to use of a continuous active motion device or continuous passive motion device for 7 days, beginning on postoperative day 1.³⁴ No difference was identified between groups in knee range of motion or pain at postoperative day 7. Patients in the continuous active motion group demonstrated a significant improvement in joint position sense (measured by passive angle reproduction) relative to the continuous passive motion group at postoperative day 7, with a between-group difference of 2.2 degrees.

Rotator Cuff Repair

Du Plessis et al (2011) published a systematic review of continuous passive motion following rotator cuff repair.³⁵ Three RCTs were included, though meta-analysis could not be conducted due to heterogeneity across trials. Two of the RCTs, by Lastayo et al (1998) and Raab et al (1996) are discussed below.^{36,37} The third trial was a German-language report by Michael et al (2005) that found a significant reduction of 12 days in the time to reach 90 abduction compared with the PT control group, with no significant difference in pain between the 2 groups.³⁸

The trial by Lastayo et al (1998) randomized 31 patients undergoing rotator cuff repair to a 4-week home program of continuous passive motion (average, 3 hours/day) or to manual passive elevation and rotation exercises.³⁶ No significant difference in outcomes was observed between the 2 approaches. Previously, Raab et al (1996) had randomized 26 patients to postoperative PT alone or to PT plus continuous passive motion.³⁷ Patients were evaluated with preoperative and 3 month postoperative shoulder scores that included pain, function, muscle strength, and range of motion. A statistically significant improvement was found in range of motion for those receiving continuous passive motion, although there was no significant improvement in overall shoulder score between groups. Both of these RCTs were likely under powered to show differences on important clinical outcomes.

Garofalo et al (2010) reported on a randomized trial assessing the effects of continuous passive motion after rotator cuff repair.³⁹ During weeks 1 to 4 post surgery, all 100 patients underwent passive self-assisted range of motion exercise, with half of the patients also receiving continuous passive motion for 4, 30-minute sessions per day. The physical therapist-supervised exercises included pendulum movements and progressive passive abduction, forward flexion, and external rotation. When patients were not exercising, the shoulder was immobilized in a sling brace. From weeks 5 to 28 post surgery, all patients underwent the same PT protocol. Visual analog scale ratings for pain were measured at 2.5, 6, and 12 months by an independent examiner. Between groups, visual analog scale ratings were slightly better for patients who received continuous passive motion at 2.5-month follow-up (7.5 vs. 9.1) but not at the 6-month (0.5 vs. 0.6) or 12-month (0.2 vs. 0.2) assessments, all respectively. Range of motion was significantly better in the group receiving continuous passive motion versus those who did not at 2.5-month follow-up

(e.g., forward flexion, 133.0° vs. 120.7°) and 6 months (158.1° vs. 151.7°) but not at 12 months (165.2° vs. 158.0°), all respectively.

Subsection Summary: Rotator Cuff Repair

Three RCTs of continuous passive motion following rotator cuff surgery were identified in the English-language literature. Two of these trials reported short-term improvements in range of motion for patients undergoing continuous passive motion, and one reported a short-term reduction in pain. None reported long-term improvements or benefits in functional status. Therefore, the clinical significance of the short-term improvements reported is uncertain. In addition, there is uncertainty about the optimal PT regimen after shoulder surgery, so the optimal comparator for continuous passive motion is not clear.

Hip Osteoarthritis

One older pilot study (1999) examined the use of continuous passive motion in patients with hip osteoarthritis in the absence of surgical intervention.⁴⁰ In this uncontrolled study, continuous passive motion was used for 1.2 to 7.6 hours daily during the 12-week trial. While improvements were noted in patients’ pain assessments, a controlled trial is needed to validate this treatment effect, particularly compared with a program of regular walking.

Femoral Fracture

Olasinde et al (2023) reported the results of a randomized trial that compared continuous passive motion to PT in patients who underwent retrograde femoral nailing for femoral fracture.⁴¹ The 88 participants were randomized to continuous passive motion or conventional PT, each for 2 hours daily. Knee stiffness at weeks 1, 2, and 6 were significantly lower among patients who received continuous passive motion compared to patients who received conventional PT (all p<.0001). Pain scores (measured by visual analogue scale) were significantly lower for the first 7 days in the continuous passive motion group, and total arc of motion gained postoperatively was also significantly larger at postoperative weeks 1, 2, and 6 (all p<.05). Interpretation of these results is limited because the duration of the intervention was not clearly stated.

ADHESIVE CAPSULITIS OF THE SHOULDER

Systematic Review

Baradaran et al (2023) conducted a systematic review of continuous passive motion compared to PT in patients with primary adhesive capsulitis (Tables 2 through 4).⁴² A total of 5 studies were included in the meta-analysis, but the conclusions were limited by heterogeneity. The authors concluded that continuous passive motion may be slightly effective in the short-term, but that long-term efficacy is still unknown.

Table 2. Trials/Studies Included in Systematic Review and Meta-Analysis

Study	Baradaran et al (2023) ⁴² ,
Azizi et al (2018)	●
Ekim et al (2016)	●
Chung et al (2015)	●

Study	Baradaran et al (2023)^{42,}
Chen et al (2009)	●
Dundar et al (2009)	●

Table 3. Systematic Review and Meta-Analysis Characteristics

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Baradaran et al (2023) ^{42,}	2009-2018	5	224	16-80	RCT	4-24 weeks

Table 4. Systematic Review and Meta-Analysis Results

Study	Pain at movement	Pain at rest	SPADI score	Constant functional shoulder score
Baradaran et al (2023) ^{42,}				
Total N	114	114	178	128
Pooled effect (95% CI)	Short-term ¹ : -1.277 (-2.146 to -0.407) Long-term ² : -1.222 (-2.224 to -0.220)	Short-term ¹ : -0.872 (-1.784 to 0.040) Long-term ² : -0.816 (-1.704 to 0.073)	Short-term ¹ : -5.196 (-12.995 to 2.602) Long-term ² : -4.561 (-12.976 to 3.855)	Short-term ¹ : 4.117 (-1.622 to 9.857) Long-term ² : 4.790 (0.376 to 9.204)
p	Short-term ¹ : .004 Long-term ² : .017	Short-term ¹ : .061 Long-term ² : .072	Short-term ¹ : .192 Long-term ² : .288	Short-term ¹ : .160 Long-term ² : .033

CI: confidence interval; SPADI: shoulder pain and disability index.

¹Short-term was defined as outcomes at week 4.

² Long-term was defined as outcomes at week 12, or pooled outcomes from weeks 8 and 24 (depending on the study).

Randomized Controlled Trials

Dundar et al (2009) compared continuous passive motion with PT in a randomized trial of 57 patients with adhesive capsulitis (frozen shoulder).⁴³ Continuous passive motion or PT was provided for 1 hour a day (5 days/week) for 4 weeks. Pain and function levels were similar in the 2 groups at baseline, with visual analog scale scores for pain ranging from 5.44 (at rest) to 6.34 (with movement). Assessments at baseline, 4, and 12 weeks showed reductions in pain and improvements in function levels for both groups. However, continuous passive motion resulted in greater pain reduction than PT (at rest, 47% vs. 25%; with movement, 35% vs. 21%; at night, 36% vs. 19%, all respectively). There were no differences between groups in range of motion or function. This trial provided modest support for the inclusion of continuous passive motion in a PT regimen for this patient population.

An RCT by Ekim et al (2016) compared continuous passive motion (n=20) with PT (n=21) for the treatment of adhesive capsulitis in patients who had diabetes.⁴⁴ Continuous passive motion or PT

was provided for 1 hour a day (5 days/week) for 4 weeks. All patients received electrotherapy and, after the 4-week initial treatment phase, were instructed to continue with an 8-week at home exercise program. Outcome measures were pain (at rest, in motion, at night) and range of motion (active and passive). Pain decreased significantly in both treatment groups, though patients in the continuous passive motion group reported a larger improvement in pain scores than those in the PT group. Range of motion improved significantly in both treatment groups as well. Patients in the continuous passive motion group reported larger improvements in abduction and flexion measures than patients in the PT group, while external and internal rotation improvements were similar across groups.

Elbow Contracture

Postoperative management of open elbow contracture release with continuous passive motion was assessed in a matched cohort study by Lindenhovius et al (2009).⁴⁵ Sixteen patients who had used continuous passive motion after open contracture release and 16 patients who had not were matched by age, sex, diagnosis, range of motion, and radiographic appearance. Improvements in range of motion did not differ between groups at the early (range, 4 to 10 months) and the final (range, 11 to 56 months) evaluations.

Hand Repair

In 1997, the TEC Assessment reviewed a multicenter study of continuous passive motion in patients who had undergone flexor tendon repair, and found the data inadequate to permit scientific conclusions about continuous passive motion application.³

Ring et al (1998) conducted a randomized trial that examined the role of continuous passive motion in patients undergoing silicone interposition arthroplasty of the metacarpophalangeal joints secondary to rheumatoid arthritis.⁴⁶ Patients were randomized to a 6-week protocol of continuous passive motion (10 hands [40 joints]) or to a standard dynamic splint protocol (15 hands [60 joints]). The trial did not show better outcomes in the continuous passive motion group.

In 2008, a retrospective chart review compared 15 patients who had received continuous passive motion after tenolysis with 21 who did not.⁴⁷ Patients who received continuous passive motion improved total active motion by 40 (range, 137 to 177), while patients who did not improved total active motion by 32 (range, 152 to 184); this difference was not statistically significant.

Foot Repair

One study (2005) has compared continuous passive motion with immobilization following surgical treatment of idiopathic club foot in 37 infants (50 feet).⁴⁸ The infants were randomized to continuous passive motion (4 hours/day) or to casting during days 10 to 42 following surgery. Blinded analysis showed improvements in the Dimeglio Clubfoot Score with continuous passive motion (range of motion, 9.7 to 3.1) that were significantly greater than those in the control group (range of motion, 10.3 to 4.2) through 12 months (97% follow-up). Between 12 and 18 months, this trend reversed and by 48 months post surgery, there was no significant difference between groups. Another study (2007) by the same group reported low compliance with this treatment.⁴⁹

Back Pain

An RCT by Gavish et al (2015) evaluated a continuous passive motion device for treatment of chronic low back pain in 36 patients.⁵⁰ Although patients treated with the device appeared to have improved outcomes on a numeric rating scale of back pain compared with waiting-list controls, the trial had significant methodologic problems. Patients who received other treatments were excluded, a large number of subjects dropped out, and control patients did not receive any conservative management.

Section Summary: Other Musculoskeletal Conditions Requiring Physical Therapy

There is a wide range of studies assessing the use of continuous passive motion for musculoskeletal conditions other than total knee arthroplasty and knee cartilage repair. No RCTs of continuous passive motion conducted in the home setting after anterior cruciate ligament repair were identified; RCTs conducted in the immediate postoperative setting do not indicate clinical benefit with use of continuous passive motion compared to conventional PT. Three small RCTs of continuous passive motion after rotator cuff surgery showed some evidence that continuous passive motion after this shoulder surgery improved short-term pain and range of motion; however, the trials were not high-quality, and the small differences in outcomes may not be clinically important. Two trials reported short-term improvements in range of motion for patients undergoing continuous passive motion, and one reported a short-term reduction in pain. None reported long-term improvements, and there are no reported benefits in functional status. Therefore, the clinical significance of the short-term improvements reported is uncertain. In addition, there is uncertainty about the optimal PT regimen following shoulder surgery such that the optimal treatment comparator for continuous passive motion is unclear. A systematic review and two small RCTs compared continuous passive motion with conventional PT for treatment of adhesive capsulitis. The systematic review was limited by heterogeneity but concluded that continuous passive motion may be effective in the short-term. One of the trials focused on diabetic patients with adhesive capsulitis. Both reported comparable improvements in range of motion and functional ability between treatment groups. For other musculoskeletal conditions, RCTs do not exist; case series either did not show efficacy of continuous passive motion or had important methodologic flaws.

STROKE**Clinical Context and Therapy Purpose**

The purpose of continuous passive motion in the home setting in individuals with stroke is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review:

Populations

The relevant population of interest is individuals with stroke.

Interventions

The therapy being considered is continuous passive motion.

Comparators

The following therapies are currently being used for stroke: standard of care.

Outcomes

The general outcomes of interest are symptoms and functional outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Continuous passive motion has been studied as a means to aid recovery of motor skills following stroke. One study (2005) randomized 35 patients to daily sessions of use of a shoulder joint continuous passive motion device (25 minutes) or to daily group therapy sessions consisting of self-directed shoulder range of motion for poststroke rehabilitation.⁵¹ All patients also received standard poststroke therapy for 3.5 hours a day. After 20 days of therapy, there was a trend for greater shoulder joint stability in the continuous passive motion group (n=17 ; p=.06) compared with the control group (n=15). No statistically significant differences were found for measures of motor impairment. This trial had a small sample size and short follow-up period, suggesting it may have had inadequate power to detect important differences in key outcomes.

In a 2022 randomized, single-blind crossover study, 18 patients aged 20 to 79 years with mild to severe arm-hand impairment following unilateral stroke were assigned (at least 6 months post-stroke) to undergo home-based therapy sessions twice daily, 5 days per week for 4 weeks, consisting of either task-specific motor training with an occupational therapist or home-based therapy with a robotic exoskeleton system combining continuous passive motion and robot-assisted gripping exercises.⁵² All patients received standard-of-care occupational therapy and PT for 2 hours per week. Crossover occurred following a 12-week washout. Patients initially assigned to the robotic exoskeleton intervention followed by task-specific motor training experienced significantly greater improvement in wrist extension range of motion at the end of treatment compared to those who received interventions in the opposite order. Assessments of manual dexterity and motor performance of the upper extremity were significantly improved following exoskeleton therapy, whereas no significant differences in these measures were noted following task-specific motor training. A significantly greater proportion of patients reported improvements in global symptoms after exoskeleton therapy (77%) than after task-specific motor training (11%).

Section Summary: Stroke

Two small randomized trials have reported mixed results with different continuous passive motion devices in combination with PT or occupational therapy compared to PT or occupational therapy alone in patients who have experienced stroke, including a statistically non-significant trend

toward improvement for the outcome of shoulder joint stability and significant improvements in wrist extension range of motion, manual dexterity, and global symptoms related to upper extremity movement. Both trials were small and treatment lasted only 20 days in the shoulder joint study by Lynch et al.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input Range of Motion Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2016 Input

In response to requests, input was received from 2 physician specialty societies and 1 academic medical center while this policy was under review in 2016. Input considered continuous passive motion (continuous passive motion) medically necessary as an adjunct to physical therapy during the non-weight-bearing rehabilitation period following articular cartilage repair procedures of the knee. One reviewer referred to the American Academy of Orthopaedic Surgery (2015) guidelines on the surgical management of osteoarthritis of the knee, which concluded that there was strong evidence that continuous passive motion after knee arthroplasty does not improve outcomes.

2010 Input

In response to requests, input was received from 2 physician specialty societies and 5 academic medical centers while this policy was under review in 2010. Overall, input supported the use of continuous passive motion under conditions of low postoperative mobility or inability to comply with rehabilitation exercises after total knee arthroplasty or total knee arthroplasty revision or during the non-weight-bearing rehabilitation period following articular cartilage repair procedures of the knee. Support was limited for use of continuous passive motion in joints other than the knee or in situations or conditions other than those described in this evidence review.

2008 Input

In response to requests, input was received from 1 physician specialty society and 2 academic medical centers while this policy was under review in 2008. The 3 reviewers interpreted the existing literature as supporting the use of continuous passive motion for the knee for at least 7 days post operatively, whether in the hospital or home, and suggested that longer use of continuous passive motion would be warranted for special conditions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Physical Therapy Association

In 2020, the American Physical Therapy Association (APTA) published a clinical practice guideline on physical therapists' management of patients undergoing total knee arthroplasty.⁵³ The APTA identified 4 high-quality studies, 6 moderate-quality studies, and 2 low-quality studies evaluating the effect of continuous passive motion devices on knee flexion and extension range of motion and need for manipulation under anesthesia, with moderate-quality studies indicating benefit with continuous passive motion contradicted by high-quality studies indicating no significant difference. Meta-analyses did not indicate a significant impact of continuous passive motion on function or hospital length of stay. The APTA concluded that "physical therapists should NOT use CPMs [continuous passive motion devices] for patients who have undergone primary, uncomplicated TKA [total knee arthroplasty]."

American Academy of Orthopaedic Surgeons

In 2015, the American Academy of Orthopaedic Surgeons (AAOS) published evidence-based guidelines on the surgical management of osteoarthritis of the knee.⁵⁴ The AAOS identified 2 high-quality studies and 5 moderate-quality studies that evaluated the use of continuous passive motion. In one high-quality study, continuous passive motion was used for about 2 weeks after discharge. The AAOS concluded that "the combined results provide strong evidence that the surgical outcomes for those who used continuous passive motion are not better than for those who did not use continuous passive motion." The 2022 update to the AAOS guidelines, which replaces the 2015 version, does not address use of continuous passive motion.⁵⁵

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.

Table 2. Summary of Key Trials

	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05952622	Comparison of Functional and Patient-reported Outcome Using Continuous Passive Motion in Rehabilitation After Plate Osteosynthesis of Proximal Humerus Fractures	103	Dec 2023
NCT05226988	Effect of Hybrid Robot-assisted Training Using End-effector and Exoskeleton Devices in Distal Upper Extremity After Stroke: Motor Control, Motor and Daily Function, Quality of Life	70	Oct 2025
<i>Unpublished</i>			
NCT01420887	Preservation of Joint Function Using Postoperative Continuous Passive Motion (continuous passive motion): A Pilot Study	60	May 2020

NCT: national clinical trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.

Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.

CPT/HCPCS	
E0935	Continuous passive motion exercise device for use on knee only
E0936	Continuous passive motion exercise device for use other than knee
E1399	Durable medical equipment, miscellaneous

REVISIONS	
03-08-2010	<p>In Policy Section:</p> <ul style="list-style-type: none"> ▪ Revised wording in C. From: "The use of a CPM device for any primary or adjunct therapeutic application other than those listed above is considered investigational/ not medically necessary." To: "The use of a CPM device for any primary or adjunct therapeutic application other than those listed above is considered experimental / investigational." ▪ Revised wording in A., third paragraph from: "CPM is approved without review-for a rental period no longer than 21 days from date of first application." To: "If CPM is approved, allow for a rental period no longer than 21 days from date of first application." <p>In Coding Section: Added HCPCS Code: E0936 (effective 01-01-07)</p> <p>Added Rationale Section</p> <p>Updated References</p>
03-07-2012	<p>In the Policy section:</p> <ul style="list-style-type: none"> ▪ In Item A, #1, removed "TKA" at the end to read "or revision." ▪ In Item A, added "#2. Partial knee arthroplasty (27446) ▪ In Item A, added the following: <ul style="list-style-type: none"> ○ "6. Osteochondral grafting ○ 7. Treatment of osteochondritis dissecans ○ 8. Repair of tibial plateau fractures" <p>Updated the Rationale section.</p> <p>Updated the Reference section.</p>
09-17-2013	<p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Added ICD-10 Diagnosis codes (<i>Effective October 1, 2014</i>) <p>Updated Rationale section.</p> <p>Updated Reference section.</p>
09-29-2015	Policy title revised from "Continuous Passive Motion for Home Use"

REVISIONS	
	Updated Description section.
	Updated Rationale section.
	Updated References section.
09-16-2016	Updated Description section.
	<p>In Policy section:</p> <ul style="list-style-type: none"> ▪ Policy language previously stated, " A.Continuous passive motion (CPM) is considered medically necessary for the following surgeries: <ol style="list-style-type: none"> 1. Total knee arthroplasty (TKA) or revision 2. Partial knee arthroplasty 3. Anterior cruciate ligament (ACL) repair 4. Autologous chondrocyte implantation (ACI) 5. Microfracture of the knee 6. Osteochondral grafting 7. Treatment of osteochondritis dissecans 8. Repair of tibial plateau fractures Continuous passive motion (CPM) will be reviewed for medical necessity for the following surgeries: <ol style="list-style-type: none"> 1. Ankle 2. Elbow 3. Wrist 4. Hand If CPM is approved, allow for a rental period no longer than 21 days from date of first application. <p>B. Continuous passive motion (CPM) is denied not medically necessary for the following surgeries:</p> <ol style="list-style-type: none"> 1. Shoulder 2. Interphalangeal joints of lower extremities to include the great toe <p>C. The use of a CPM device for any primary or adjunct therapeutic application other than those listed above is considered experimental / investigational."</p> <ul style="list-style-type: none"> ▪ Current policy language states, " A.Use of continuous passive motion (CPM) in the home setting may be considered medically necessary as an adjunct to physical therapy in the following situations: <ol style="list-style-type: none"> 1. Under conditions of low postoperative mobility or inability to comply with rehabilitation exercises following a total knee arthroplasty (TKA) or TKA revision. This may include patients with complex regional pain syndrome (reflex sympathetic dystrophy); extensive arthrofibrosis or tendon fibrosis; or physical, mental, or behavioral inability to participate in active physical therapy. 2. During the non-weight-bearing rehabilitation period following intra-articular cartilage repair procedures of the knee (e.g., microfracture, osteochondral grafting, autologous chondrocyte implantation, treatment of osteochondritis dissecans, repair of tibial plateau fractures. <p>B. Use of CPM in the home setting for all other conditions is considered not medically necessary.</p> <p>Policy Guidelines</p> <ol style="list-style-type: none"> 1. If CPM is approved, allow for a rental period no longer than 21 days from date of first application."
	Updated Rationale section.
	Updated References section.
04-12-2017	Updated Description section.

REVISIONS	
	Updated Rationale section.
	Updated References section.
04-11-2018	Updated Description section.
	Updated Rationale section.
	In Coding section:
	▪ Removed ICD-9 codes.
	Updated References section.
04-24-2019	Updated Rationale section.
	Updated References section.
06-07-2021	Updated Description section.
	Updated Policy Guidelines
	• Removed "If CPM is approved, allow for a rental period no longer than 21 days from date of first application"
	• Added Policy Guideline 1 and 2.
	Updated Rationale section.
	Updated References section.
05-04-2022	Updated Description Section
	Updated Rationale Section
	Updated Coding Section
	▪ Converted ICD-10 Codes to ranges
	Updated References Section
04-25-2023	Updated Description Section
	Updated Rationale Section
	Updated Coding Section
	▪ Removed ICD-10 Codes
	Updated References Section
04-23-2024	Updated Description Section
	Updated Rationale Section
	Updated References Section

REFERENCES

1. O'Driscoll SW, Giori NJ. Continuous passive motion (CPM): theory and principles of clinical application. J Rehabil Res Dev. 2000; 37(2): 179-88. PMID 10850824
2. Gholson JJ, Noiseux NO, Otero JE, et al. Patient Factors Systematically Influence Hospital Length of Stay in Common Orthopaedic Procedures. Iowa Orthop J. 2017; 37: 233-237. PMID 28852363
3. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Continuous Passive Motion as an Adjunct to Physical Therapy for Joint Rehabilitation. TEC Assessments. 1997;Volume 12:Tab 20.
4. McInnes J, Larson MG, Daltroy LH, et al. A controlled evaluation of continuous passive motion in patients undergoing total knee arthroplasty. JAMA. Sep 16 1992; 268(11): 1423-8. PMID 1512910
5. Milne S, Brosseau L, Robinson V, et al. Continuous passive motion following total knee arthroplasty. Cochrane Database Syst Rev. 2003; (2): CD004260. PMID 12804511
6. Brosseau L, Milne S, Wells G, et al. Efficacy of continuous passive motion following total knee arthroplasty: a metaanalysis. J Rheumatol. Nov 2004; 31(11): 2251-64. PMID 15517640

7. Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database Syst Rev*. Feb 06 2014; 2014(2): CD004260. PMID 24500904
8. Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database Syst Rev*. Mar 17 2010; (3): CD004260. PMID 20238330
9. He ML, Xiao ZM, Lei M, et al. Continuous passive motion for preventing venous thromboembolism after total knee arthroplasty. *Cochrane Database Syst Rev*. Jul 29 2014; 2014(7): CD008207. PMID 25069620
10. Yashar AA, Venn-Watson E, Welsh T, et al. Continuous passive motion with accelerated flexion after total knee arthroplasty. *Clin Orthop Relat Res*. Dec 1997; (345): 38-43. PMID 9418619
11. MacDonald SJ, Bourne RB, Rorabeck CH, et al. Prospective randomized clinical trial of continuous passive motion after total knee arthroplasty. *Clin Orthop Relat Res*. Nov 2000; (380): 30-5. PMID 11064970
12. Pope RO, Corcoran S, McCaul K, et al. Continuous passive motion after primary total knee arthroplasty. Does it offer any benefits?. *J Bone Joint Surg Br*. Nov 1997; 79(6): 914-7. PMID 9393903
13. Kumar PJ, McPherson EJ, Dorr LD, et al. Rehabilitation after total knee arthroplasty: a comparison of 2 rehabilitation techniques. *Clin Orthop Relat Res*. Oct 1996; (331): 93-101. PMID 8895624
14. Bruun-Olsen V, Heiberg KE, Mengshoel AM. Continuous passive motion as an adjunct to active exercises in early rehabilitation following total knee arthroplasty - a randomized controlled trial. *Disabil Rehabil*. 2009; 31(4): 277-83. PMID 18608367
15. Denis M, Moffet H, Caron F, et al. Effectiveness of continuous passive motion and conventional physical therapy after total knee arthroplasty: a randomized clinical trial. *Phys Ther*. Feb 2006; 86(2): 174-85. PMID 16445331
16. Leach W, Reid J, Murphy F. Continuous passive motion following total knee replacement: a prospective randomized trial with follow-up to 1 year. *Knee Surg Sports Traumatol Arthrosc*. Oct 2006; 14(10): 922-6. PMID 16489477
17. Boese CK, Weis M, Phillips T, et al. The efficacy of continuous passive motion after total knee arthroplasty: a comparison of three protocols. *J Arthroplasty*. Jun 2014; 29(6): 1158-62. PMID 24412145
18. Herbold JA, Bonistall K, Blackburn M, et al. Randomized controlled trial of the effectiveness of continuous passive motion after total knee replacement. *Arch Phys Med Rehabil*. Jul 2014; 95(7): 1240-5. PMID 24685389
19. Chen B, Zimmerman JR, Soulen L, et al. Continuous passive motion after total knee arthroplasty: a prospective study. *Am J Phys Med Rehabil*. 2000; 79(5): 421-6. PMID 10994883
20. Herbold JA, Bonistall K, Blackburn M. Effectiveness of continuous passive motion in an inpatient rehabilitation hospital after total knee replacement: a matched cohort study. *PM R*. Oct 2012; 4(10): 719-25. PMID 22959052
21. Worland RL, Arredondo J, Angles F, et al. Home continuous passive motion machine versus professional physical therapy following total knee replacement. *J Arthroplasty*. Oct 1998; 13(7): 784-7. PMID 9802665

22. Lenssen TA, van Steyn MJ, Crijns YH, et al. Effectiveness of prolonged use of continuous passive motion (CPM), as an adjunct to physiotherapy, after total knee arthroplasty. *BMC Musculoskelet Disord*. Apr 29 2008; 9: 60. PMID 18442423
23. Browne JE, Anderson AF, Arciero R, et al. Clinical outcome of autologous chondrocyte implantation at 5 years in US subjects. *Clin Orthop Relat Res*. Jul 2005; (436): 237-45. PMID 15995447
24. Farr J. Autologous chondrocyte implantation improves patellofemoral cartilage treatment outcomes. *Clin Orthop Relat Res*. Oct 2007; 463: 187-94. PMID 17960681
25. Rosenberger RE, Gomoll AH, Bryant T, et al. Repair of large chondral defects of the knee with autologous chondrocyte implantation in patients 45 years or older. *Am J Sports Med*. Dec 2008; 36(12): 2336-44. PMID 18725654
26. Nugent-Derfus GE, Takara T, O'Neill JK, et al. Continuous passive motion applied to whole joints stimulates chondrocyte biosynthesis of PRG4. *Osteoarthritis Cartilage*. May 2007; 15(5): 566-74. PMID 17157538
27. Salter RB. The biologic concept of continuous passive motion of synovial joints. The first 18 years of basic research and its clinical application. *Clin Orthop Relat Res*. May 1989; (242): 12-25. PMID 2650945
28. Fazalare JA, Griesser MJ, Siston RA, et al. The use of continuous passive motion following knee cartilage defect surgery: a systematic review. *Orthopedics*. Dec 01 2010; 33(12): 878. PMID 21162503
29. Howard JS, Mattacola CG, Romine SE, et al. Continuous Passive Motion, Early Weight Bearing, and Active Motion following Knee Articular Cartilage Repair: Evidence for Clinical Practice. *Cartilage*. Oct 2010; 1(4): 276-86. PMID 26069559
30. Hill AD, Palmer MJ, Tanner SL, et al. Use of Continuous Passive Motion in the Postoperative Treatment of Intra-Articular Knee Fractures. *J Bone Joint Surg Am*. Jul 16 2014; 96(14): e118. PMID 25031380
31. Wright RW, Preston E, Fleming BC, et al. A systematic review of anterior cruciate ligament reconstruction rehabilitation: part I: continuous passive motion, early weight bearing, postoperative bracing, and home-based rehabilitation. *J Knee Surg*. Jul 2008; 21(3): 217-24. PMID 18686484
32. Culvenor AG, Girdwood MA, Juhl CB, et al. Rehabilitation after anterior cruciate ligament and meniscal injuries: a best-evidence synthesis of systematic reviews for the OPTIKNEE consensus. *Br J Sports Med*. Dec 2022; 56(24): 1445-1453. PMID 35768181
33. Gatewood CT, Tran AA, Dragoo JL. The efficacy of post-operative devices following knee arthroscopic surgery: a systematic review. *Knee Surg Sports Traumatol Arthrosc*. Feb 2017; 25(2): 501-516. PMID 27695905
34. Friemert B, Bach C, Schwarz W, et al. Benefits of active motion for joint position sense. *Knee Surg Sports Traumatol Arthrosc*. Jun 2006; 14(6): 564-70. PMID 16328464
35. Du Plessis M, Eksteen E, Jenneker A, et al. The effectiveness of continuous passive motion on range of motion, pain and muscle strength following rotator cuff repair: a systematic review. *Clin Rehabil*. Apr 2011; 25(4): 291-302. PMID 20943710
36. Lastayo PC, Wright T, Jaffe R, et al. Continuous passive motion after repair of the rotator cuff. A prospective outcome study. *J Bone Joint Surg Am*. Jul 1998; 80(7): 1002-11. PMID 9698005
37. Raab MG, Rzeszutko D, O'Connor W, et al. Early results of continuous passive motion after rotator cuff repair: a prospective, randomized, blinded, controlled study. *Am J Orthop (Belle Mead NJ)*. Mar 1996; 25(3): 214-20. PMID 8775698

38. Michael JW, König DP, Imhoff AB, et al. [Efficiency of a postoperative treatment after rotator cuff repair with a continuous passive motion device (CPM)]. *Z Orthop Ihre Grenzgeb.* 2005; 143(4): 438-45. PMID 16118760
39. Garofalo R, Conti M, Notarnicola A, et al. Effects of one-month continuous passive motion after arthroscopic rotator cuff repair: results at 1-year follow-up of a prospective randomized study. *Musculoskelet Surg.* May 2010; 94 Suppl 1: S79-83. PMID 20383685
40. Simkin PA, de Lateur BJ, Alquist AD, et al. Continuous passive motion for osteoarthritis of the hip: a pilot study. *J Rheumatol.* Sep 1999; 26(9): 1987-91. PMID 10493681
41. Olasinde AA, Olisa O, Muhumuza J, et al. Early outcome measurement of the effectiveness of conventional physical therapy versus continuous passive motion in knee function following retrograde femoral nailing-a prospective randomized controlled trial. *Int Orthop.* Aug 2023; 47(8): 2085-2093. PMID 37269402
42. Baradaran A, Ebrahimzadeh MH, Sabzevari S, et al. Is there any advantage between using continuous passive motion and conventional physical therapy in patients with primary adhesive capsulitis?: A systematic review and meta-analysis. *J Bodyw Mov Ther.* Oct 2023; 36: 133-141. PMID 37949549
43. Dundar U, Toktas H, Cakir T, et al. Continuous passive motion provides good pain control in patients with adhesive capsulitis. *Int J Rehabil Res.* Sep 2009; 32(3): 193-8. PMID 19011582
44. Ekim AA, İnal EE, Gönüllü E, et al. Continuous passive motion in adhesive capsulitis patients with diabetes mellitus: A randomized controlled trial. *J Back Musculoskelet Rehabil.* Nov 21 2016; 29(4): 779-786. PMID 27002662
45. Lindenhovius AL, van de Luitgaarden K, Ring D, et al. Open elbow contracture release: postoperative management with and without continuous passive motion. *J Hand Surg Am.* 2009; 34(5): 858-65. PMID 19362791
46. Ring D, Simmons BP, Hayes M. Continuous passive motion following metacarpophalangeal joint arthroplasty. *J Hand Surg Am.* May 1998; 23(3): 505-11. PMID 9620192
47. Schwartz DA, Chafetz R. Continuous passive motion after tenolysis in hand therapy patients: a retrospective study. *J Hand Ther.* 2008; 21(3): 261-6; quiz 267. PMID 18652971
48. Zeifang F, Carstens C, Schneider S, et al. Continuous passive motion versus immobilisation in a cast after surgical treatment of idiopathic club foot in infants: a prospective, blinded, randomised, clinical study. *J Bone Joint Surg Br.* Dec 2005; 87(12): 1663-5. PMID 16326882
49. Kasten P, Geiger F, Zeifang F, et al. Compliance with continuous passive movement is low after surgical treatment of idiopathic club foot in infants: a prospective, double-blinded clinical study. *J Bone Joint Surg Br.* Mar 2007; 89(3): 375-7. PMID 17356153
50. Gavish L, Barzilay Y, Koren C, et al. Novel continuous passive motion device for self-treatment of chronic lower back pain: a randomised controlled study. *Physiotherapy.* Mar 2015; 101(1): 75-81. PMID 25280603
51. Lynch D, Ferraro M, Krol J, et al. Continuous passive motion improves shoulder joint integrity following stroke. *Clin Rehabil.* Sep 2005; 19(6): 594-9. PMID 16180594
52. Kuo LC, Yang KC, Lin YC, et al. Internet of Things (IoT) Enables Robot-Assisted Therapy as a Home Program for Training Upper Limb Functions in Chronic Stroke: A Randomized Control Crossover Study. *Arch Phys Med Rehabil.* Mar 2023; 104(3): 363-371. PMID 36122608

53. Jette DU, Hunter SJ, Burkett L, et al. Physical Therapist Management of Total Knee Arthroplasty. *Phys Ther*. Aug 31 2020; 100(9): 1603-1631. PMID 32542403
54. American Academy of Orthopaedic Surgeons. Surgical management of osteoarthritis of the knee: Evidence-based clinical practice guideline. Rosemont, IL: AAOS; 2015.
55. American Academy of Orthopaedic Surgeons. Surgical Management of Osteoarthritis of the Knee Evidence Based Clinical Practice Guideline. Published December 2, 2022. Accessed January 29, 2024.
56. Center for Medicare & Medicaid. National Coverage Decision (NCD) for Durable Medical Equipment Reference List (280.1). 2023; <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=190&ncdver=3&bc=0>. Accessed January 29, 2024.

OTHER REFERENCES

1. Blue Cross and Blue Shield of Kansas Orthopedic Liaison Committee, January 10, 2007 (see Blue Cross and Blue Shield of Kansas Newsletter, Blue Shield Report. MAC-01-07).
2. Blue Cross and Blue Shield of Kansas Medical Advisory Committee meeting, April 19, 2007 (see Blue Cross and Blue Shield of Kansas Newsletter, Blue Shield Report. MAC-01-07).
3. Orthopedic Liaison Committee January 5, 2007 and Medical Advisory Committee April 19, 2007.
4. Blue Cross and Blue Shield of Kansas Orthopedic Liaison Committee, March 2009; June 2016.