

Medical Policy



Title: Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

| | |
|---|--|
| Professional / Institutional | |
| Original Effective Date: August 23, 2013 | |
| Latest Review Date: January 1, 2025 | |
| Current Effective Date: February 15, 2018 | |

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

| Populations | Interventions | Comparators | Outcomes |
|---|--|--|--|
| Individuals: <ul style="list-style-type: none"> With suspected inherited motor and sensory peripheral neuropathy | Interventions of interest are: <ul style="list-style-type: none"> Testing for genes associated with inherited peripheral neuropathies | Comparators of interest are: <ul style="list-style-type: none"> Clinical management without genetic testing | Relevant outcomes include: <ul style="list-style-type: none"> Test validity Symptoms Change in disease status |

DESCRIPTION

The inherited peripheral neuropathies are a heterogeneous group of diseases that may be inherited in an autosomal dominant, autosomal recessive, or X-linked dominant manner. These diseases can generally be diagnosed based on clinical presentation, nerve conduction studies, and family history. Genetic testing has been used to diagnose specific inherited peripheral neuropathies.

OBJECTIVE

The objective of this evidence review is to evaluate whether genetic testing in individuals with suspected inherited motor and sensory peripheral neuropathy improves health outcomes.

BACKGROUND**Inherited Peripheral Neuropathies**

Inherited peripheral neuropathies are a clinically and genetically heterogeneous group of disorders. The estimated prevalence in aggregate is 1 in 2500 persons, making inherited peripheral neuropathies the most common inherited neuromuscular disease.¹

Peripheral neuropathies can be subdivided into 2 major categories: primary axonopathies and primary myelinopathies, depending on which portion of the nerve fiber is affected. The further anatomic classification includes fiber type (e.g., motor vs. sensory, large vs. small) and gross distribution of the nerves affected (e.g., symmetry, length-dependency).

Inherited peripheral neuropathies are divided into hereditary motor and sensory neuropathies, hereditary neuropathy with liability to pressure palsies (HNPP), and other miscellaneous, rare types (e.g., hereditary brachial plexopathy, hereditary sensory, autonomic neuropathies). Other hereditary metabolic disorders, such as Friedreich ataxia, Refsum disease, and Krabbe disease, may be associated with motor and/or sensory neuropathies but typically have other predominating symptoms. This evidence review focuses on hereditary motor and sensory neuropathies and HNPP.

A genetic etiology of peripheral neuropathy is typically suggested by generalized polyneuropathy, family history, lack of positive sensory symptoms, early age of onset, symmetry, associated skeletal abnormalities, and very slowly progressive clinical course.² A family history of at least 3 generations with details on health issues, the cause of death, and age at death should be collected.

CHARCOT-MARIE-TOOTH DISEASE**Hereditary Motor and Sensory Neuropathies**

Most inherited polyneuropathies were originally described clinically as variants of Charcot-Marie-Tooth (CMT) disease. The clinical phenotype of CMT is highly variable, ranging from minimal neurologic findings to the classic picture with pes cavus and "stork legs" to a severe polyneuropathy with respiratory failure.³ CMT disease is genetically and clinically heterogeneous. Variants in more than 30 genes and more than 44 different genetic loci have been associated with inherited neuropathies.⁴ Also, different pathogenic variants in a single gene can

lead to different inherited neuropathy phenotypes and inheritance patterns. A 2016 cross-sectional study of 520 children and adolescents with CMT found variability in CMT-related symptoms across the 5 most commonly represented subtypes.⁵

CMT subtypes are characterized by variants in 1 of several myelin genes, which lead to abnormalities in myelin structure, function, or upkeep. There are 7 subtypes of CMT, with type 1 and 2 representing the most common hereditary peripheral neuropathies.

Most cases of CMT are autosomal dominant, although autosomal recessive and X-linked dominant forms exist. Most cases are CMT type 1 (approximately 40% to 50% of all CMT cases, with 78% to 80% of those due to *PMP22* variants).⁶ CMT type 2 is associated with 10% to 15% of CMT cases, with 20% of those due to *MFN2* variants.

A summary of the molecular genetics of CMT is outlined in Table 1.

Table 1. Molecular Genetics of CMT Variants

| Locus | Gene | Protein Product | Prevalence (if known) |
|--------------|---------------|---|------------------------------|
| CMT type 1 | | | |
| CMT1A | <i>PMP22</i> | Peripheral myelin protein 22 | 50% of CMT1 |
| CMT1B | <i>MPZ</i> | Myelin P0 protein | 25% of CMT1 |
| CMT1C | <i>LITAF</i> | Lipopolysaccharide-induced tumor necrosis factor- α factor | |
| CMT1D | <i>EGR2</i> | Early growth response protein 2 | |
| CMT1E | <i>PMP22</i> | Peripheral myelin protein 22 (sequence changes) | |
| CMT1F/2E | <i>NEFL</i> | Neurofilament light polypeptide | |
| CMT1G | <i>PMP2</i> | Peripheral myelin protein 2 | |
| CMT type 2 | | | |
| CMT2A1 | <i>KIF1B</i> | Kinesin-like protein KIF1B | |
| CMT2A2A/B | <i>MFN2</i> | Mitofusin-2 | |
| CMT2B | <i>RAB7A</i> | Ras-related protein Rab-7 | |
| CMT2B1 | <i>LMNA</i> | Lamin A/C | |
| CMT2B2 | <i>PNKP</i> | | |
| CMT2C | <i>TRPV4</i> | Transient receptor potential cation channel subfamily V member 4 | |
| CMT2D | <i>GARS1</i> | Glycyl-tRNA synthetase | |
| CMT2F | <i>HSPB1</i> | Heat-shock protein beta-1 | |
| CMT2G | <i>LRSAM1</i> | E3 ubiquitin-protein ligase LRSAM1 | |
| CMT2H/2K | <i>GDAP1</i> | Ganglioside-induced differentiation-associated protein 1 | |

| Locus | Gene | Protein Product | Prevalence (if known) |
|--------------|----------------|---|------------------------------|
| CMT2I/ J | <i>MPZ</i> | Myelin P0 protein | |
| CMT2L | <i>HSPB8</i> | Heat-shock protein beta-8 | |
| CMT2M | <i>DNM2</i> | Dynamin 2 | |
| CMT2N | <i>AARS1</i> | Alanyl-tRNA synthetase, cytoplasmic | |
| CMT2O | <i>DYNC1H1</i> | Cytoplasmic dynein 1 heavy chain 1 | |
| CMT2P | <i>LRSAM1</i> | E3 ubiquitin-protein ligase LRSAM1 | |
| CMT2Q | <i>DHTKD1</i> | Dehydrogenase E1 And Transketolase Domain Containing 1 | |
| CMT2R | <i>TRIM2</i> | Tripartite Motif Containing 2 | |
| CMT2S | <i>IGHMBP2</i> | DNA-binding protein SMUBP-2 | |
| CMT2T | <i>MME</i> | Membrane Metalloendopeptidase | |
| CMT2U | <i>MARS1</i> | Methionine-tRNA ligase, cytoplasmic | |
| CMT2V | <i>NAGLU</i> | N-Acetyl-Alpha-Glucosaminidase | |
| CMT2W | <i>HARS1</i> | Histidyl-TRNA Synthetase 1 | |
| CMT2X | <i>SPG11</i> | Spastic paraplegia 11 | |
| CMT2Y | <i>VCP</i> | Valosin Containing Protein | |
| CMT2Z | <i>MORC2</i> | Microrchidia Family CW-Type Zinc Finger 2 | |
| CMT type 4 | | | |
| CMT4A | <i>GDAP1</i> | Ganglioside-induced differentiation-associated protein 1 | |
| CMT4B1 | <i>MTMR2</i> | Myotubularin-related protein 2 | |
| CMT4B2 | <i>SBF2</i> | Myotubularin-related protein 13 | |
| CMT4B3 | <i>SBF1</i> | SET Binding Factor 1 | |
| CMT4C | <i>SH3TC2</i> | SH3 domain and tetratricopeptide repeats-containing protein 2 | |
| CMT4D | <i>NDRG1</i> | Protein NDRG1 | |
| CMT4E | <i>EGR2</i> | Early growth response protein 2 | |
| CMT4F | <i>PRX</i> | Periaxin | |
| CMT4H | <i>FGD4</i> | FYVE, RhoGEF, and PH domain-containing protein 4 | |
| CMT4J | <i>FIG4</i> | Phosphatidylinositol 3, 5-biphosphate | |
| X-linked CMT | | | |
| CMTX3 | <i>Xq26</i> | Unknown | |
| CMTX4 | <i>AIFM1</i> | Apoptosis-inducing factor 1 | |

| Locus | Gene | Protein Product | Prevalence (if known) |
|-------|--------------|---|-----------------------|
| CMTX5 | <i>PRPS1</i> | Ribose-phosphate pyrophosphokinase 1 | |
| CMTX6 | <i>PDK3</i> | Pyruvate dehydrogenase kinase isoform 3 | |

Adapted from Bird (2022).⁶
CMT: Charcot-Marie-Tooth.

CMT Type 1

CMT1 is an autosomal dominant, demyelinating peripheral neuropathy characterized by distal muscle weakness and atrophy, sensory loss, and slow nerve conduction velocity. It is usually slowly progressive and often associated with pes cavus foot deformity, bilateral foot drop, and palpably enlarged nerves, especially the ulnar nerve at the olecranon groove and the greater auricular nerve. Affected people usually become symptomatic between ages 5 and 25 years, and their lifespan is not shortened. Less than 5% of people become wheelchair-dependent. CMT1 is inherited in an autosomal dominant manner. The CMT1 subtypes (CMT 1A-E) are separated by molecular findings and are often clinically indistinguishable. CMT1A accounts for 70% to 80% of all CMT1, and about two-thirds of probands with CMT1A have inherited the disease-causing variant, and about one-third have CMT1A as the result of a de novo variant.

CMT1A involves duplication of the *PMP22* gene. *PMP22* encodes an integral membrane protein, peripheral membrane protein 22, which is a major component of myelin in the peripheral nervous system. The phenotypes associated with this disease arise because of abnormal *PMP22* gene dosage effects.⁷ Two normal alleles represent the normal wild-type condition. Four normal alleles (as in the homozygous CMT1A duplication) result in the most severe phenotype, whereas 3 normal alleles (as in the heterozygous CMT1A duplication) cause a less severe phenotype.⁶

CMT Type 2

CMT2 is a non-demyelinating (axonal) peripheral neuropathy characterized by distal muscle weakness and atrophy, mild sensory loss, and normal or near-normal nerve conduction velocities. Clinically, CMT2 is similar to CMT1, although typically less severe.⁶ The subtypes of CMT2 are similar clinically and distinguished only by molecular genetic findings. CMT2B1, CMT2B2, and CMT2H/K are inherited in an autosomal recessive manner; all other subtypes of CMT2 are inherited in an autosomal dominant manner. The most common subtype of CMT2 is CMT2A, which accounts for approximately 20% of CMT2 cases and is associated with variants in the *MFN2* gene.

X-Linked CMT

CMT X type 1 is characterized by a moderate-to-severe motor and sensory neuropathy in affected males and mild to no symptoms in carrier females.⁸ Sensorineural deafness and central nervous system symptoms also occur in some families. CMT X type 1 is inherited in an X-linked dominant manner. Molecular genetic testing of *GJB1* (Cx32), which is available on a clinical basis, detects about 90% of cases of CMT X type 1.

CMT Type 4

CMT type 4 is a form of hereditary motor and sensory neuropathy that is inherited in an autosomal recessive fashion and occurs secondary to myelinopathy or axonopathy. It occurs more rarely than the other forms of CMT neuropathy, but some forms may be rapidly progressive and/or associated with severe weakness.

Hereditary Neuropathy with Liability to Pressure Palsies

The largest proportion of CMT1 cases are due to variants in *PMP22*. In HNPP (also called tomaculous neuropathy), inadequate production of *PMP22* causes nerves to be more susceptible to trauma or minor compression or entrapment. Patients with HNPP rarely present symptoms before the second or third decade of life. However, some have reported presentation as early as birth or as late as the seventh decade of life.⁹ The prevalence is estimated at 16 persons per 100,000, although some authors have indicated a potential for underdiagnosis of the disease.⁹ An estimated 50% of carriers are asymptomatic and do not display abnormal neurologic findings on clinical examination.¹⁰ HNPP is characterized by repeated focal pressure neuropathies such as carpal tunnel syndrome and peroneal palsy with foot drop and episodes of numbness, muscular weakness, atrophy, and palsies due to minor compression or trauma to the peripheral nerves. The disease is benign with complete recovery occurring within a period of days to months in most cases, although an estimated 15% of patients have residual weakness following an episode.¹⁰ Poor recovery usually involves a history of prolonged pressure on a nerve, but, in these cases, the remaining symptoms are typically mild.

PMP22 is the only gene for which a variant is known to cause HNPP. A large deletion occurs in approximately 80% of patients, and the remaining 20% of patients have single nucleotide variants (SNVs) and small deletions in the *PMP22* gene. One normal allele (due to a 17p11.2 deletion) results in HNPP and a mild phenotype. SNVs in *PMP22* have been associated with a variable spectrum of HNPP phenotypes ranging from mild symptoms to representing a more severe, CMT1-like syndrome.¹¹ Studies have also reported that the SNV frequency may vary considerably by ethnicity.¹² About 10% to 15% of variant carriers remain clinically asymptomatic, suggesting incomplete penetrance.¹³

Treatment

Currently, there is no therapy to slow the progression of neuropathy for inherited peripheral neuropathies. A 2015 systematic review of exercise therapies for CMT including 9 studies described in 11 articles reported significant improvements with functional activities and physiological adaptations with exercise.¹⁴ Supportive treatment, if necessary, is generally provided by a multidisciplinary team including neurologists, physiatrists, orthopedic surgeons, and physical and occupational therapists. Treatment choices are limited to physical therapy, the use of orthotics, surgical treatment for skeletal or soft tissue abnormalities, and drug treatment for pain.¹⁵ Avoidance of obesity and drugs associated with nerve damage (e.g., vincristine, paclitaxel, cisplatin, isoniazid, nitrofurantoin) is recommended for patients with CMT.⁶

Supportive treatment for HNPP can include transient bracing (e.g., wrist splint or ankle-foot orthosis), which may become permanent in some cases of foot drop.¹⁶ Prevention of HNPP manifestations can be accomplished by wearing protective padding (e.g., elbow or knee pads) to prevent trauma to nerves during activity. Some have reported that vincristine should also be avoided in HNPP patients.^{6,16} Ascorbic acid has been investigated as a treatment for CMT1A based on animal models, but a 2013 trial in humans did not demonstrate significant clinical benefit.¹⁷ Attarian et al (2014) reported results of an exploratory phase 2 randomized, double-blind, placebo-controlled trial of PXT3003, a low-dose combination of 3 approved compounds (baclofen, naltrexone, sorbitol) in 80 adults with CMT1A.¹⁸ The trial demonstrated the safety and tolerability of the drug. Mandel et al (2015) included this randomized controlled trial and 3 other trials (1 of ascorbic acid, 2 of PXT3003) in a meta-analysis.¹⁹

Molecular Genetic Testing

Multiple laboratories offer individual variant testing for genes involved in hereditary sensory and motor neuropathies, which would typically involve sequencing analysis via Sanger sequencing or next-generation sequencing followed by deletion/duplication analysis (i.e., with array comparative genomic hybridization) to detect large deletions or duplications. For the detection of variants in *MFN2*, whole gene or select exome sequence analysis is typically used to identify SNVs, in addition to or followed by deletion or duplication analysis for the detection of large deletions or duplications.

Aretz et al (2010) reported a general estimation of the clinical sensitivity of CMT variant testing for hereditary motor and sensory neuropathy and HNPP using a variety of analytic methods (multiplex ligation-dependent probe amplification, multiplex amplicon quantification, quantitative polymerase chain reaction, Southern blot, fluorescence in-situ hybridization, pulsed-field gel electrophoresis, denaturing high-performance liquid chromatography, high-resolution melting, restriction analysis, direct sequencing).²⁰ The clinical sensitivity (i.e., the proportion of positive tests if the disease is present) for the detection of deletions/duplications or mutations to *PMP22* was about 50% and 1%, respectively, for single nucleotide variants. The clinical specificity (i.e., the proportion of negative tests if the disease is not present) was nearly 100%.

A number of genetic panel tests for the assessment of peripheral neuropathies are commercially available. For example, GeneDx (Gaithersburg, MD) offers an Axonal CMT panel, which uses next-generation sequencing and exon array comparative genomic hybridization. The genes tested include *AARS, AIFM1, BSCL2, DNAJB2, DNM2, DYNC1H1, GAN, GARS, GDAP1, GJB1, GNB4, HARS, HINT1, HSPB1, HSPB8, IGHMBP2, INF2, KIF5A, LMNA, LRSAM1, MFN2, MME, MORC2, MPZ, NEFL, PLEKHG5, PRPS1, RAB7A, SLC12A6, TRIM2, TRPV4, and YARS*.²¹ InterGenetics (Athens, Greece) offers a next-generation sequencing panel for neuropathy that includes 42 genes involved in CMT, along with other hereditary neuropathies. Fulgent Clinical Diagnostics Lab offers a broader next-generation sequencing panel for CMT that includes 48 genes associated with CMT and other neuropathies and myopathies.

REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Genetic testing for the diagnosis of inherited peripheral neuropathies is available under the auspices of CLIA. Laboratories that offer laboratory-developed tests must be licensed by CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

POLICY

- A. Genetic testing is considered **medically necessary** when the diagnosis of an inherited peripheral motor or sensory neuropathy is suspected due to signs and/or symptoms, but a definitive diagnosis cannot be made without genetic testing.
- B. Genetic testing for an inherited peripheral neuropathy is considered **experimental / investigational** for all other indications.

POLICY GUIDELINES

- A. This policy addresses the hereditary motor and sensory peripheral neuropathies, of which peripheral neuropathy is the primary clinical manifestation. A number of other hereditary disorders may have neuropathy as an associated finding but typically have other central nervous system or other systemic findings. Examples include Refsum disease, various lysosomal storage diseases, and mitochondrial disorders.

- B. **Genetics Nomenclature Update**

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Variome Project, the Human Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology - "pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign" - to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

| Previous | Updated | Definition |
|----------|----------------------------|---|
| Mutation | Disease-associated variant | Disease-associated change in the DNA sequence |
| | Variant | Change in the DNA sequence |
| | Familial variant | Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives |

Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification

| Variant Classification | Definition |
|-----------------------------------|--|
| Pathogenic | Disease-causing change in the DNA sequence |
| Likely pathogenic | Likely disease-causing change in the DNA sequence |
| Variant of uncertain significance | Change in DNA sequence with uncertain effects on disease |
| Likely benign | Likely benign change in the DNA sequence |
| Benign | Benign change in the DNA sequence |

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

C. Genetic Counseling

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods

Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was conducted through November 17, 2023. (see Appendix Table 1 for genetic testing categories).

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

TESTING FOR GENES ASSOCIATED WITH INHERITED PERIPHERAL NEUROPATHIES

Clinical Context and Test Purpose

The purpose of testing for variants associated with hereditary motor and sensory neuropathies in individuals with suspected inherited peripheral neuropathy is to make a diagnosis of an inherited peripheral neuropathy or to inform the prognosis of inherited peripheral neuropathy.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with suspected inherited peripheral neuropathy who present with sensory, motor, or mixed findings, and sometimes with other findings. Charcot-Marie-Tooth (CMT) disease is clinically heterogeneous.

Interventions

The relevant intervention of interest is testing for variants associated with CMT, by deletion or duplication analysis, usually by multiplex ligation-dependent probe amplification, and gene sequencing, usually by next-generation sequencing.

Genetic counseling is particularly important for CMT given the extreme genetic heterogeneity of the disorder.

Comparators

The relevant comparator of interest is a clinical diagnosis of an inherited peripheral neuropathy made using a combination of clinical features, family pedigree, and characteristic nerve conduction velocity/electromyography studies. However, subtypes of CMT are defined based on their genotype.

Outcomes

The general outcomes of interest are test validity, symptoms, and change in disease status. Beneficial outcomes resulting from a true test include avoiding potentially harmful therapies. Harmful outcomes resulting from a false-positive test include potentially unneeded treatments due to misidentified individuals.

Testing can be conducted during diagnostic evaluation, and follow-up should be continued for years after diagnosis.

Study Selection Criteria

For the evaluation of clinical validity, studies that meet the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Review of Evidence

England et al (2009) reported on the role of laboratory and genetic tests in the evaluation of distal symmetric polyneuropathies and concluded that genetic testing is established as useful for the accurate diagnosis and classification of hereditary polyneuropathies in patients with a cryptogenic polyneuropathy who exhibit a classical hereditary neuropathy phenotype.³ Six studies included in the review showed that when the test for *CMT1A* duplication is restricted to patients with clinically probable CMT1 (i.e., autosomal dominant, primary demyelinating polyneuropathy), the yield is 54% to 80%, compared with testing a cohort of patients suspected of having any variety of hereditary peripheral neuropathies, where the yield is only 25% to 59% (average, 43%).

Sequential Testing

Given the genetic complexity of CMT, many commercial and private laboratories evaluate CMT with a testing algorithm based on patients' presenting characteristics. For the evaluation of the clinical validity of genetic testing for CMT, we included studies that evaluated patients with clinically suspected CMT who were evaluated with a genetic testing algorithm that was described in the study.

Saporta et al (2011) reported results from genetic testing of 1024 patients with clinically suspected CMT who were evaluated at a single institution's CMT clinic from 1997 to 2009.⁴ Patients who were included were considered to have CMT if they had sensorimotor peripheral neuropathy and a family history of a similar condition. Patients without a family history of neuropathy were considered to have CMT if their medical history, neurophysiologic testing, and neurologic examination were typical for CMT1, CMT2, CMTX, or CMT4. Seven hundred eighty-seven patients were diagnosed with CMT; of those, 527 (67%) had a specific genetic diagnosis as a result of their visit. Genetic testing decisions were left up to the treating clinician, and the authors noted that decisions about which genes to test changed during the study. Most (98.2%) of those with clinically diagnosed CMT1 had a genetic diagnosis, and of all patients with a genetic diagnosis, most (80.8%) had clinically diagnosed CMT1. The authors characterized several clinical phenotypes of CMT based on clinical presentation and physiologic testing.

Rudnik-Schoneborn et al (2016) reported on results from genetic testing of 1206 index patients and 124 affected relatives who underwent genetic testing at a single reference laboratory from 2001 to 2012.²² Patients were referred by neurologic or genetic centers throughout Germany, and were grouped by age at onset (early infantile [<2 years], childhood [2 to 10 years], juvenile [10 to 20 years], adult [20 to 50 years], late adult [>50 years]), and by electroneurographic

findings. Molecular genetic methods changed over the course of the study, and testing was tiered by patient features and family history. Of the 674 index patients with a demyelinating CMT phenotype on nerve conduction studies, 343 (51%) had a genetic diagnosis; of the 340 index patients with an axonal CMT phenotype, 45 (13%) had a genetic diagnosis; and of the 192 with hereditary neuropathy with liability to pressure palsies (HNPP), 67 (35%) had a genetic diagnosis. The most common genetic diagnoses differed by nerve conduction phenotype: of the 429 patients genetically identified with demyelinating CMT (index and secondary), 89.3% were detected with *PMP22* deletion or duplication (74.8%), *GJB1/Cx32* (8.9%), or *MPZ/PO* (5.6%) variant analysis. In contrast, of the 57 patients genetically identified with axonal CMT (index and secondary), 84.3% were detected with *GJB1/Cx32* (42.1%), *MFN2* (33.3%), or *MPZ/PO* (8.8%) variant analysis.

In an earlier study, Gess et al (2013) reported on sequential genetic testing for CMT-related genes from 776 patients at a single center for suspected inherited peripheral neuropathies from 2004 to 2012.²³ Most patients (n=624) were treated in the same center. The test strategy varied based on electrophysiologic data and family history. The testing yield was 66% (233/355) in patients with CMT1, 35% (53/151) in patients with CMT2, and 64% (53/83) in patients with HNPP. Duplications on chromosome 17 were the most common variants in CMT1 (77%), followed by *GJB1* (13%) and *MPZ* (8%) variants among those with positive genetic tests. For CMT2 patients, *GJB2* (30%) and *MFN2* (23%) variants were most common among those with positive genetic tests.

Ostern et al (2013) reported on a retrospective analysis of cases of CMT diagnostic testing referred to a single reference laboratory in Norway from 2004 to 2010.²⁴ Genetic testing was stratified based on clinical information supplied on patient requisition forms based on age of onset of symptoms, prior testing, results from motor nerve conduction velocity, and patterns of inheritance. The study sample included 435 index cases of a total of 549 CMT cases tested (other tests were for at-risk family members or other reasons). Patients were grouped based on whether they had symptoms of polyneuropathy, classical CMT, with or without additional symptoms or changes in imaging or had atypical features or the physician suspected an alternative diagnosis. Among the cases tested, 72 (16.6%) were found to be variant-positive, all of whom had symptoms of CMT. Most (69/72 [95.8%]) of the positive molecular genetic findings were *PMP22* region duplications or sequence variants in *MPZ*, *GJB1*, or *MFN2* genes.

Murphy et al (2012) reported on the yield of sequential testing for CMT-related gene variants from 1607 patients with testing sent to a single center.²⁵ Of the 916 patients seen in the authors' clinic, 601 (65.6%) had a primary inherited neuropathy, including 425 with CMT and 46 with HNPP. Of the 425 with a clinical diagnosis of CMT, 240 had CMT1 (56.5%), and 115 (27.1%) had CMT2. Of those with CMT, 266 (62.6%) of 425 received a genetic diagnosis, most frequently (92%) with a variant in 1 of 4 genes (*PMP22* duplication, and *GJB1*, *MPZ*, and *MFN2*).

Uchôa Cavalcanti (2021) reported on results from genetic testing of 503 patients (94 families and 192 unrelated individuals) who underwent testing in a Brazilian neuromuscular outpatient clinic from 2015 to 2020.²⁶ The diagnosis of CMT was established based on the presence of slowly progressive, motor and sensory neuropathy, independent of any family history. Patients were assessed utilizing clinical and neurophysiological data along with targeted gene panel sequencing. Among the 503 patients, a genetic diagnosis was reported in 394 patients (77 families and 120 unrelated individuals). The following confirmed genetic diagnoses were identified: demyelinating

CMT (n=317), intermediate CMT (n=34), and axonal CMT (n=43). The genetic diagnosis rate in probands was 68.9% (197/286). The most common causative genes were *PMP22* duplication *GJB1*, *MFN2*, *GDAP1*, *MPZ*, *PMP22* point mutation, *NEFL*, *SBF2*, and *SH3TC2*.

In addition to sequential testing algorithms, some studies were reported on the yield of multigene testing panels, most often using next-generation sequencing methods. Studies with populations of suspected inherited motor or sensory neuropathy that have reported on next-generation sequencing panel test results are summarized in Table 2.

Table 2. Summary of Genetic Panel Tests in Charcot-Marie-Tooth

| Study | N | Population | Test | Diagnostic Yield (NGS Panel) | VUS (NGS Panel) |
|---|----------------------|---|--|--|-------------------------------|
| Antoniadi et al (2015) ²⁷ , | 448 | Suspected inherited peripheral neuropathy, with supportive NCV, some with negative testing for <i>PMP22</i> | 56-gene NGS panel | 137 (31%) patients (31 genes) | NR |
| DiVincenzo et al (2014) ²⁸ , | 17,377; 503 with NGS | Suspected peripheral neuropathy, referred to a central laboratory | 14-gene NGS panel and <i>PMP22</i> del/dup by MLPA | 95 (18.9%) patients (8 genes) | 38 (7.5%) patients (11 genes) |
| Volodarsky et al (2021) ²⁹ , | 2517 | Suspected diagnosis of CMT, referred to a molecular genetics laboratory | 34-gene NGS panel | 440 (17.5%) patients; 6 genes constituted 80% of the overall results | NR |

CMT: Charcot-Marie-Tooth; del/dup: deletion/duplication; MLPA: multiplex ligation-dependent amplification; NCV: nerve conduction velocity; NGS: next-generation sequencing; NR: not reported; VUS: variant of uncertain significance.

Genotype-Phenotype Correlations

There is significant clinical variability within and across subtypes of CMT. Therefore, some studies have evaluated genotype-phenotype correlations within CMT cases. For example, Sanmaneechai et al (2015) characterized genotype-phenotype correlations in patients with CMT1B regarding *MPZ* variants in a cohort of 103 patients from 71 families.³⁰ Patients underwent standardized clinical assessments and clinical electrophysiology. There were 47 different *MPZ* variants and 3 characteristic ages of onset: infantile (age range, 0 to 5 years), childhood (age range, 6 to 20 years), and adult (age range, ≥21 years). Specific variants were clustered by age group, with only 2 variants found in more than 1 age group.

Karadima et al (2015) investigated the association between *PMP22* variants and clinical phenotype in 100 Greek patients referred for genetic testing for HNPP.³¹ In the 92 index cases, the frequency of *PMP22* deletions was 47.8%, and the frequency of *PMP22* micro-variants was 2.2%. Variant-negative patients were more likely to have an atypical phenotype (41%), absent family history (96%), and nerve conduction study findings not fulfilling HNPP criteria (80.5%).

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

The clinical utility of genetic testing for hereditary peripheral neuropathies depends on how the results can be used to improve patient management. Published data for the clinical utility of genetic testing for inherited peripheral neuropathies is lacking.

The diagnosis of an inherited peripheral neuropathy can generally be made clinically. However, when the diagnosis cannot be made clinically, a genetic diagnosis may add incremental value. A diagnosis of an inherited peripheral neuropathy is important to direct therapy, regarding early referrals to physical therapy and avoidance of potentially toxic medications. Some specific medications for CMT are under investigation, but their use is not well-established. There are significant differences in prognosis for different forms of CMT, although whether a different prognosis leads to choices in therapy that lead to different outcomes is uncertain. In some cases, a genetic diagnosis of an inherited peripheral neuropathy may have the potential to avoid other diagnostic tests.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

No direct evidence for improved outcomes with the use of genetic testing for hereditary motor and sensory peripheral neuropathies and HNPP was identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity.

There is evidence from observational studies to support the use of genetic testing to establish a diagnosis in cases of suspected inherited motor or sensory neuropathy when a diagnosis cannot be made by other methods and, in turn, to initiate supportive therapies.

Section Summary: Testing for Genes Associated with Inherited Peripheral Neuropathies

A relatively large body of literature, primarily from retrospective, single-center reference labs in which patients with suspected CMT have been tested, addressed clinical validity. The testing yield is reasonably high, particularly when patients are selected based on clinical phenotype.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Neurology

In 2009, the American Academy of Neurology (AAN) and 2 other specialty societies published an evidence-based, tiered approach for the evaluation of distal symmetric polyneuropathy and suspected hereditary neuropathies, which concluded the following (see Table 3).³

Table 3. Recommendations on Distal Symmetric Polyneuropathy and Suspected Hereditary Neuropathies

| Recommendation | LOE ^a |
|--|------------------|
| "Genetic testing is established as useful for the accurate diagnosis and classification of hereditary neuropathies" | A |
| "Genetic testing may be considered in patients with cryptogenic polyneuropathy who exhibit a hereditary neuropathy phenotype" | C |
| "Initial genetic testing should be guided by the clinical phenotype, inheritance pattern, and electrodiagnostic features and should focus on the most common abnormalities which are CMT1A duplication/HNPP deletion, Cx32 (GJB1), and MFN2 screening" | |
| "There is insufficient evidence to determine the usefulness of routine genetic testing in patients with cryptogenic polyneuropathy who do not exhibit a hereditary neuropathy phenotype" | U |

CMT: Charcot-Marie-Tooth; HNPP: hereditary neuropathy with liability to pressure palsies; LOE: level of evidence.

^a Grade A: established as effective, ineffective, or harmful for the given condition in the specified population; grade C: possibly effective, ineffective, or harmful for the given condition in the specified population; grade U: data inadequate or conflicting; given current knowledge.

The AAN website indicates the recommendations were reaffirmed on January 22, 2022, and indicated an update is in progress.

American Academy of Family Physicians

In 2020, the American Academy of Family Physicians recommended genetic testing for a patient with suspected peripheral neuropathy, if basic blood tests are negative, electrodiagnostic studies suggest an axonal etiology and diseases such as diabetes, toxic medications, thyroid disease, vitamin deficiency, and vasculitis can be ruled out.³²

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 4.

Table 4. Summary of Key Trials

| NCT No. | Trial Name | Planned Enrollment | Completion Date |
|----------------|--|---------------------------|------------------------|
| <i>Ongoing</i> | | | |
| NCT01193075 | Natural History Evaluation of Charcot Marie Tooth Disease (CMT) Type (CMT1B), 2A (CMT2A), 4A (CMT4A), 4C (CMT4C), and Others | 5000 | December 2024 |
| NCT01193088 | Genetics of Charcot Marie Tooth Disease (CMT) - Modifiers of CMT1A, New Causes of CMT | 1050 | December 2024 |

NCT: national clinical trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.

Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.

| CPT/HCPCS | |
|------------------|--|
| 81324 | PMP22 (peripheral myelin protein 22) (e.g., Charcot-Marie-Tooth, hereditary neuropathy with liability to pressure palsies) gene analysis; duplication/deletion analysis |
| 81325 | PMP22 (peripheral myelin protein 22) (e.g., Charcot-Marie-Tooth, hereditary neuropathy with liability to pressure palsies) gene analysis; full sequence analysis |
| 81326 | PMP22 (peripheral myelin protein 22) (e.g., Charcot-Marie-Tooth, hereditary neuropathy with liability to pressure palsies) gene analysis; known familial variant |
| 81403 | Molecular pathology procedure, Level 4 (e.g., analysis of single exon by DNA sequence analysis, analysis of >10 amplicons using multiplex PCR in 2 or more independent reactions, mutation scanning or duplication/ deletion variants of 2-5 exons) |
| 81404 | Molecular pathology procedure, Level 5 (e.g., analysis of 2-5 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 6-10 exons, or characterization of a dynamic mutation disorder/triplet repeat by Southern blot analysis) |
| 81405 | Molecular pathology procedure, Level 6 (e.g., analysis of 6-10 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 11-25 exons, regionally targeted cytogenomic array analysis) |
| 81406 | Molecular pathology procedure, Level 7 (e.g., analysis of 11-25 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 26-50 exons, cytogenomic array analysis for neoplasia) |
| 81448 | Hereditary peripheral neuropathies (e.g. Charcot-Marie-Tooth, spastic paraplegia), genomic sequence analysis panel, must include sequencing of at least 5 peripheral neuropathy-related genes (e.g., BSCL2, GJB1, MFN2, MPZ, REEP1, SPAST, SPG11, SPTLC1) |
| 83884 | Neurofilament light chain (eff. 01-01-2025) |
| 81479 | Unlisted molecular pathology procedure |

| REVISIONS | |
|------------------|---|
| 08-23-2013 | Policy posted to the bcbsks.com web site on 07-24-2013 for an effective date of 08-23-2013. |
| 10-06-2015 | Description section added. |
| | In Policy section: <ul style="list-style-type: none"> ▪ Policy guidelines added containing clarifying information about the variety of neuropathies and genetic counseling. |
| | Rationale section updated |
| | In Coding section: <ul style="list-style-type: none"> ▪ Added CPT Codes: 81404, 81405, 81406, 81479 ▪ Coding notations updated |
| | References updated |
| 02-15-2018 | Policy published 12-29-2017. Policy effective 02-15-2018. |
| | Description section updated |
| | In Policy section: <ul style="list-style-type: none"> ▪ In Item A revised policy position from experimental / investigational to medically necessary to read "Genetic testing is considered medically necessary when the diagnosis of an inherited peripheral motor or sensory neuropathy is suspected due to signs and/or symptoms but a definitive diagnosis cannot be made without genetic testing." |
| | Rationale section updated |
| | In Coding section: <ul style="list-style-type: none"> ▪ Added CPT Code: 81448 (Effective 01-01-2018) ▪ Added ICD-10 Codes: G60.0, G60.8, G60.9 ▪ Coding notations updated |
| | References updated |
| 04-10-2019 | Description section updated |
| | In Policy section: <ul style="list-style-type: none"> ▪ Policy Guidelines updated |
| | Rationale section updated |
| | In Coding section: <ul style="list-style-type: none"> ▪ Added CPT Code: 81403 |
| | References updated |
| 02-18-2021 | Description section updated |
| | Rationale section updated |
| | References updated |
| 03-08-2022 | Updated Description Section |
| | Updated Policy guidelines <ul style="list-style-type: none"> ▪ Added "Genetics Nomenclature Update" Section |
| | Updated Rationale Section |
| | Updated Coding Section <ul style="list-style-type: none"> ▪ Removed Coding bullets <ul style="list-style-type: none"> ○ There is specific CPT coding for genetic testing for <i>PMP22</i> deletions and duplications, full sequencing, and familial variant testing: 81324, 81325, 81326, 81448. ○ CPT Tier 2 code 81403 includes the following test- ○ <i>GJB1</i> (<i>gap junction protein, beta 1</i>) (e.g., Charcot-Marie-Tooth X-linked), full gene sequence. ○ CPT Tier 2 code 81404 includes the following tests- ○ <i>EGR2</i> (<i>early growth response 2</i>) (e.g., Charcot-Marie-Tooth), full gene sequence ○ <i>HSPB1</i> (<i>heat shock 27kDa protein 1</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence |

| REVISIONS | |
|------------------|--|
| | <ul style="list-style-type: none"> ○ <i>LITAF</i> (<i>lipopolysaccharide-induced TNF factor</i>) (e.g., Charcot-Marie-Tooth), full gene sequence ○ CPT Tier 2 code 81405 includes the following tests– ○ <i>GDAP1</i> (<i>ganglioside-induced differentiation-associated protein 1</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence. ○ <i>MPZ</i> (<i>myelin protein zero</i>) (e.g., Charcot-Marie-Tooth), full gene sequence ○ <i>NEFL</i> (<i>neurofilament, light polypeptide</i>) (e.g., Charcot-Marie-Tooth), full gene sequence ○ <i>PRX</i> (<i>periaxin</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence ○ <i>RAB7A</i> (<i>RAB7A, member RAS oncogene family</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence. ○ CPT Tier 2 code 81406 includes the following tests– ○ <i>FIG4</i> (<i>FIG4 homolog, SAC1 lipid phosphatase domain containing [S. cerevisiae]</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence ○ <i>GARS</i> (<i>glycyl-tRNA synthetase</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence ○ <i>LMNA</i> (<i>lamin A/C</i>) (e.g., Emery-Dreifuss muscular dystrophy [EDMD1, 2 and 3] limb-girdle muscular dystrophy [LGMD] type 1B, dilated cardiomyopathy [CMD1A], familial partial lipodystrophy [FPLD2]), full gene sequence ○ <i>MFN2</i> (<i>mitofusin 2</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence. ○ <i>SH3TC2</i> (<i>SH3 domain and tetratricopeptide repeats 2</i>) (e.g., Charcot-Marie-Tooth disease), full gene sequence ○ For the other genes listed above, there is no specific CPT listing of the test and the unlisted molecular pathology code 81479 would be reported. |
| | Updated References Section |
| 02-28-2023 | Updated Description Section |
| | Updated Rationale Section |
| | Updated References Section |
| | Removed Appendix Section |
| 03-12-2024 | Updated Description Section |
| | Updated Rationale Section |
| | Updated Coding Section <ul style="list-style-type: none"> ▪ Removed ICD-10 Codes |
| | Updated References Section |
| 01-01-2025 | Updated Coding Section <ul style="list-style-type: none"> ▪ Added: 83884 (eff. 01-01-2025) |

REFERENCES

1. Burgunder JM, Schöls L, Baets J, et al. EFNS guidelines for the molecular diagnosis of neurogenetic disorders: motoneuron, peripheral nerve and muscle disorders. *Eur J Neurol.* Feb 2011; 18(2): 207-217. PMID 20500522
2. Alport AR, Sander HW. Clinical approach to peripheral neuropathy: anatomic localization and diagnostic testing. *Continuum (Minneapolis, Minn).* Feb 2012; 18(1): 13-38. PMID 22810068

3. England JD, Gronseth GS, Franklin G, et al. Practice Parameter: evaluation of distal symmetric polyneuropathy: role of laboratory and genetic testing (an evidence-based review). Report of the American Academy of Neurology, American Association of Neuromuscular and Electrodiagnostic Medicine, and American Academy of Physical Medicine and Rehabilitation. *Neurology*. Jan 13 2009; 72(2): 185-92. PMID 19056666
4. Saporta AS, Sottile SL, Miller LJ, et al. Charcot-Marie-Tooth disease subtypes and genetic testing strategies. *Ann Neurol*. Jan 2011; 69(1): 22-33. PMID 21280073
5. Cornett KM, Menezes MP, Bray P, et al. Phenotypic Variability of Childhood Charcot-Marie-Tooth Disease. *JAMA Neurol*. Jun 01 2016; 73(6): 645-51. PMID 27043305
6. Bird TD. Charcot-Marie-Tooth Hereditary Neuropathy Overview. In: Adam MP, Everman DB, Mirzaa GM, et al., editors. *GeneReviews*. Seattle, WA: University of Washington; 2023.
7. Stankiewicz P, Lupski JR. The genomic basis of disease, mechanisms and assays for genomic disorders. *Genome Dyn*. 2006; 1: 1-16. PMID 18724050
8. Abrams CK. GJB1 Disorders: Charcot Marie Tooth Neuropathy (CMT1X) and Central Nervous System Phenotypes. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2020.
9. Meretoja P, Silander K, Kalimo H, et al. Epidemiology of hereditary neuropathy with liability to pressure palsies (HNPP) in south western Finland. *Neuromuscul Disord*. Dec 1997; 7(8): 529-32. PMID 9447611
10. Celik Y, Kilincer C, Hamamcioğlu MK, et al. Hereditary neuropathy with liability to pressure palsies in a Turkish patient (HNPP): a rare cause of entrapment neuropathies in young adults. *Turk Neurosurg*. Jan 2008; 18(1): 82-4. PMID 18382985
11. Taioli F, Cabrini I, Cavallaro T, et al. Inherited demyelinating neuropathies with micromutations of peripheral myelin protein 22 gene. *Brain*. Feb 2011; 134(Pt 2): 608-17. PMID 21252112
12. Bissar-Tadmouri N, Parman Y, Boutrand L, et al. Mutational analysis and genotype/phenotype correlation in Turkish Charcot-Marie-Tooth Type 1 and HNPP patients. *Clin Genet*. Nov 2000; 58(5): 396-402. PMID 11140841
13. Dubourg O, Mouton P, Brice A, et al. Guidelines for diagnosis of hereditary neuropathy with liability to pressure palsies. *Neuromuscul Disord*. Mar 2000; 10(3): 206-8. PMID 10734269
14. Sman AD, Hackett D, Fiatarone Singh M, et al. Systematic review of exercise for Charcot-Marie-Tooth disease. *J Peripher Nerv Syst*. Dec 2015; 20(4): 347-62. PMID 26010435
15. Pareyson D, Marchesi C. Natural history and treatment of peripheral inherited neuropathies. *Adv Exp Med Biol*. 2009; 652: 207-24. PMID 20225028
16. Chrestian N. Hereditary neuropathy with liability to pressure palsies. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *Gene Reviews*. Seattle, WA: University of Washington; 2020.
17. Lewis RA, McDermott MP, Herrmann DN, et al. High-dosage ascorbic acid treatment in Charcot-Marie-Tooth disease type 1A: results of a randomized, double-masked, controlled trial. *JAMA Neurol*. Aug 2013; 70(8): 981-7. PMID 23797954
18. Attarian S, Vallat JM, Magy L, et al. An exploratory randomised double-blind and placebo-controlled phase 2 study of a combination of baclofen, naltrexone and sorbitol (PXT3003) in patients with Charcot-Marie-Tooth disease type 1A. *Orphanet J Rare Dis*. Dec 18 2014; 9: 199. PMID 25519680

19. Mandel J, Bertrand V, Lehert P, et al. A meta-analysis of randomized double-blind clinical trials in CMT1A to assess the change from baseline in CMTNS and ONLS scales after one year of treatment. *Orphanet J Rare Dis.* Jun 13 2015; 10: 74. PMID 26070802
20. Aretz S, Rautenstrauss B, Timmerman V. Clinical utility gene card for: HMSN/HNPP HMSN types 1, 2, 3, 6 (CMT1,2,4, DSN, CHN, GAN, CCFDN, HNA); HNPP. *Eur J Hum Genet.* Sep 2010; 18(9). PMID 20512157
21. GeneDx. Charcot-Marie-Tooth Panel. <https://www.genedx.com/tests/detail/charcot-marie-tooth-panel-888> Accessed November 17, 2023.
22. Rudnik-Schöneborn S, Tölle D, Senderek J, et al. Diagnostic algorithms in Charcot-Marie-Tooth neuropathies: experiences from a German genetic laboratory on the basis of 1206 index patients. *Clin Genet.* Jan 2016; 89(1): 34-43. PMID 25850958
23. Gess B, Schirmacher A, Boentert M, et al. Charcot-Marie-Tooth disease: frequency of genetic subtypes in a German neuromuscular center population. *Neuromuscul Disord.* Aug 2013; 23(8): 647-51. PMID 23743332
24. Østern R, Fagerheim T, Hjeltnes H, et al. Diagnostic laboratory testing for Charcot Marie Tooth disease (CMT): the spectrum of gene defects in Norwegian patients with CMT and its implications for future genetic test strategies. *BMC Med Genet.* Sep 21 2013; 14: 94. PMID 24053775
25. Murphy SM, Laura M, Fawcett K, et al. Charcot-Marie-Tooth disease: frequency of genetic subtypes and guidelines for genetic testing. *J Neurol Neurosurg Psychiatry.* Jul 2012; 83(7): 706-10. PMID 22577229
26. Uchôa Cavalcanti EB, Santos SCL, Martins CES, et al. Charcot-Marie-Tooth disease: Genetic profile of patients from a large Brazilian neuromuscular reference center. *J Peripher Nerv Syst.* Sep 2021; 26(3): 290-297. PMID 34190362
27. Antoniadis T, Buxton C, Dennis G, et al. Application of targeted multi-gene panel testing for the diagnosis of inherited peripheral neuropathy provides a high diagnostic yield with unexpected phenotype-genotype variability. *BMC Med Genet.* Sep 21 2015; 16: 84. PMID 26392352
28. DiVincenzo C, Elzinga CD, Medeiros AC, et al. The allelic spectrum of Charcot-Marie-Tooth disease in over 17,000 individuals with neuropathy. *Mol Genet Genomic Med.* Nov 2014; 2(6): 522-9. PMID 25614874
29. Volodarsky M, Kerkhof J, Stuart A, et al. Comprehensive genetic sequence and copy number analysis for Charcot-Marie-Tooth disease in a Canadian cohort of 2517 patients. *J Med Genet.* Apr 2021; 58(4): 284-288. PMID 32376792
30. Sanmaneechai O, Feely S, Scherer SS, et al. Genotype-phenotype characteristics and baseline natural history of heritable neuropathies caused by mutations in the MPZ gene. *Brain.* Nov 2015; 138(Pt 11): 3180-92. PMID 26310628
31. Karadima G, Koutsis G, Raftopoulou M, et al. Mutational analysis of Greek patients with suspected hereditary neuropathy with liability to pressure palsies (HNPP): a 15-year experience. *J Peripher Nerv Syst.* Jun 2015; 20(2): 79-85. PMID 26110377
32. Castelli G, Desai KM, Cantone RE. Peripheral Neuropathy: Evaluation and Differential Diagnosis. *Am Fam Physician.* Dec 15 2020; 102(12): 732-739. PMID 33320513